

## SUICIDE AND ATTEMPTED SUICIDE

RMA ID Number	Reference List for RMA250-6 as at August 2018
------------------	---

T
Mi A, Fruchter E, Mann JJ, et al (2013). Differentiating army suicide attempters from psychologically treated and untreated soldiers: a demographic, psychological and stress-reaction characterization. J Affect Disord, 150: 300-5.
Aardema H, Meertens JH, Ligtenberg JJ, et al (2008). Organophosphorus pesticide poisoning: cases and developments. The Netherlands Journal of Medicine, 66(4): 149-53.
Adam I, Elbashir MI (2004). Suicide after treatment of chloroquine- resistant falciparum malaria with quinine. Saudi Med J, 25(2): 248-9.
Adam KS (1985). Attempted suicide. Psychiatr Clin N Am, 8(2): 183-201.
Adams J, Adams M (1996). The association among negative life events, perceived problem solving alternatives, depression, and suicidal ideation in adolescent psychiatric patients. J Child Psychol, 37(6): 715-20.
Adapted from Sadock (1980). From the legal definition of rape in the United States (Brownmiller 1975). Kaplans Comprehensive Textbook of Psychiatry, Third Edition, 1803.
Adshead Surg Lt S (2014). Clinical research. The adverse effects of mefloquine in deployed military personnel. J R Nav Med Serv, 100(3): 232-7.
Agerbo E (2005). Midlife suicide risk, partner's psychiatric illness, spouse and child bereavement by suicide or other modes of death: a gender specific study. J Epidemiol Community Health, 59: 407-12.
Agoub M, Moussaoui D, Kadri N (2006). Assessment of suicide in a Moroccan metropolitan area. Journal of Affective Disorders, 90: 223-6.
Ahmed K, Mohan RA, Bhugra D (2007). Self-harm in South Asian women: A literature review informed approach to assessment and formulation. Am J Psychother, 61(1): 71-81.
Ahn E, Shin DW, Cho S-I, et al (2010). Suicide rates and risk factors among Korean cancer patients, 1993-2005. Cancer Epidemiol Biomarkers Prev, 19(8): 2097-105.
Akyuz G, Sar V, Kugu N, et al (2005). Reported childhood trauma, attempted suicide and self-mutilative behavior among women in the general population. Eur Psychiatry, 20: 268-73.
Ali AK, Heran BS, Etminan M (2015). Persistent sexual dysfunction and suicidal ideation in young men treated with low-dose finasteride: a pharmacovigilance study. Pharmacotherapy, 35(7): 687-95.
Allebeck P, Bolund C (1991). Suicides and suicide attempts in cancer patients. Psychol Med, 21(4): 979-84.
Allen JP, Cross G, Swanner J (2005). Suicide in the army: a review of current information. Military Medicine, 170: 580-4.
Allgulander C (1994). Suicide and mortality patterns in anxiety neurosis and depressive neurosis. Arch Gen Psychiatry, 51: 708-12.

	Allgulander C, Lavori PW (1993). Causes of death among 936 elderly
4181	patients with 'pure' anxiety neurosis in Stockholm County, Sweden, and
1101	in patients with depressive neurosis or both diagnoses. Comprehensive
	Psychiatry, 34(5): 299-302.
4212	Allgulander C, Lavori PW (1991). Excess mortality among 3302 patients
	with 'pure' anxiety neurosis. Arch Gen Psychiatry, 48: 599-602.
4=40	American Psychiatric Association (1994). Diagnostic and statistical
1746	manual of mental disorders. Personality Disorders, 4th Edition. American
	Psychiatric Association, Washington DC.
70700	American Psychiatric Association (2013). Diagnostic and statistical
70783	manual of mental disorders. DSM-5, 5th Edition, American Psychiatric
	Publishing, Inc.
77004	Amitai M, Apter A (2012). Social aspects of suicidal behavior and
77691	prevention in early life: a review. Int J Environ Res Public Health, 9: 985-
	94.
4740	Anderson, DM, Keith J, Novak PD (Lexicographers) (1994). Dorland's
1749	Illustrated Medical Dictionary 28th Edition, 1601. W B Saunders
	Philadelphia.  Anderson, DM, Keith J, Novak PD (Lexicographers) (1994). Dorland's
1750	Illustrated Medical Dictionary, 28th Edition,: 1232. W B Saunders
1730	Philadelphia.
	Andrews JA, Lewinsohn PM (1992). Suicidal attempts among older
4206	adolescents: Prevalence and co-occurrence with psychiatric disorders. J
1200	Am Acad Child Adolesc Psychiatry, 31(4): 655-62.
	Ang RP, Ooi YP (2004). Impact of gender and parents' marital status on
52467	adolescents' suicidal ideation. IJSP, 50(4): 351-60.
	Angst J, Degonda M, Ernst C (1992). The Zurich Study: XV. Suicide
4211	attempts in a cohort from age 20 to 30. Eur Arch Psychiatry Clin
	Neurosci, 242: 135-41.
E 4000	Anon (2009). Rimonabant: depression and suicide. Prescrire
54266	International, 18(99): 24.
53889	Anonymous (2008). Suicidal behaviour and self harm. The facts.
33003	Retrieved 28 August 2009, from http://www.sane.org
	Antretter E, Dunkel D, Haring C, et al (2008). The factorial structure of
55083	the Suicide Intent Scale: a comparative study in clinical samples from 11
	European regions. Int J Methods Psychiatr Res, 17(2): 63-79.
4400	Appleby L (1994). Panic and suicidal behaviour: Risk of self-harm in
4183	patients who complain of panic. British Journal of Psychiatry, 164(164):
	719-21.
1740	Appleby L (1994). Panic and suicidal behaviour: Risk of self-harm in
	patients who complain of panic. Br J Psychiatry, 164(6): 719-21.  Arie M, Haruvi-Catalan L, Apter A (2005). Personality and suicidal
54725	behavior in adolescence. Clinical Neuropsychiatry, 2(1): 37-47.
	Ariga M, Uehara T, Takeuchi K, et al (2008). Trauma exposure and
86458	posttraumatic stress disorder in delinquent female adolescents. Journal
30 100	of Child Psychology and Psychiatry, 49(1): 79-87.
	Atigari OV, Hogan C, Healy D (2013). Doxycycline and suicidality. BMJ
77551	Case Rep, doi:10.1136/bcr-2013-200723.
50400	Australian Bureau of Statistics (2007). National survey of mental health
52106	and wellbeing: summary of results. 4326.0: 1-98.
	Australian Institute of Health and Welfare (2017). Incidence of suicide
85057	among service and ex-serving Australian Defence Force personnel 2001-
	2015: in brief summary report. Cat. No. PHE 213. Australian Institute of
	Health and Welfare.
	Australian Institute of Health and Welfare (2016). Incidence of suicide
81438	among serving and ex-serving Australian Defence Force personnel2001-
	2014.

85058	Australian Institute of Health and Welfare (2017). Incidence of suicide in
	serving and ex-serving Australia Defence Force personnel: detailed
	analysis 2001-2015. Cat. No. PHE 218. Canberra. Australian Institute of Health and Welfare.
19155	Australian Institute of Health and Welfare [AIHW] (2000). Suicide in
	Vietnam Veterans' children. Morbidity of Vietnam Veterans,
	Supplementary Report No. 1, No. PHE 25. AIHW, Canberra.
54763	Averbach IJ (1973). History of psychiatric disorder and risk of academic
34703	failure in freshmen. J Am Coll Health Assoc, 21(3): 252-6.
	Ayalon L, Mackin S, Arean PA, et al (2007). The role of cognitive
54501	functioning and distress in suicidal ideation in older adults. J Am Geriatr
	Soc, 55: 1090-4.
54000	Azorin JM, Kaladjian A, Adida M, et al (2009). Risk factors associated
51968	with lifetime suicide attempts in bipolar I patients: findings from a French
	national cohort. Comprehensive Psychiatry, 50: 115-20.
77000	Baghaei P, Tabarsi P, Dorriz D, et al (2011). Adverse effects of
77829	multidrug-resistant tuberculosis treatment with a standardized regimen: a
	report from Iran. Am J Ther, 18(2): e29-34.
77930	Bahmanyar S, Sparen P, Mittendorfer Rutz E, et al (2009). Risk of suicide among operated and non-operated patients hospitalised for
11930	peptic ulcers. J Epidemiol Comm Health, 63(12): 1016-21.
	Bailey JM (1999). [Comment] Homosexuality and mental illness. Arch
53683	Gen Psychiatry, 56: 883-4. Comment on ID: 53682.
	Bang SM, Lee JH, Yoon SS, et al (2006). A multicenter retrospective
54044	analysis of adverse events in Korean patients using bortezomib for
0.01.	multiple myeloma. Int J Hematol, 83: 309-13.
	Bangs ME, Tauscher-Wisniewski S, Polzer J, et al (2008). Meta-analysis
54292	of suicide-related behavior events in patients treated with atomoxetine. J
	Am Acad Child Adolesc Psychiatry, 47(2): 209-18.
	Bangs ME, Wietecha LA, Wang S, et al (2014). Meta-analysis of suicide-
77794	related behavior of ideation in child, adolescent, and adult patients
11194	treated with atomoxetine. Journal of Child and Adolescent
	Psychopharmacology, 24(8): 426-34.
53501	Barbour V, Clark J, Peiperl L, et al (2009). [Comment] Media portrayals of
00001	suicide. PLoS Med, 6(3): e1000051. Comment on ID: 53445.
52456	Barbour V, Clark J, Peiperl L, et al (2009). [Comment] Media portrayals of
52.50	suicide. PLoS Med, 6(3): e1000051. Comment on ID: 53445.
5.4000	Barbui C, Esposito E, Cipriani A (2009). Selective serotonin reuptake
54323	inhibitors and risk of suicide: a systematic review of observational
	studies. CMAJ, 180(3): 291-7.
86455	Bardon C, Mishara BL (2015). Systematic review of the impact of suicides and other critical incidents on railway personnel. Suicide and
00400	Life-Threatening Behavior, 45(6): 720-31.
	Barnhorst A (2015). Amphetamine-related psychiatric disorders
78097	Retrieved 29 April 2016, from
10091	http://emedicine.medscape.com/article/289973-overview
	Basham C, Denneson LM, Millet L, et al (2011). Characteristics and VA
	health care utilization of U.S. veterans who completed suicide in Oregon
79641	between 2000 and 2005. Suicide & Life-Threatening Behaviour, 41(3):
	287-96.
	Bastia BK, Kar N (2009). A psychological autopsy study of suicidal
54825	hanging from Cuttack, India: focus on stressful life situations. Arch
	Suicide Res, 13: 100-4.
	Bastia BK, Kar N (2009). A psychological autopsy study of suicidal
54721	hanging from Cuttack, India: focus on stressful life situations. Arch
	Suicide Res, 13: 100-4.

79839	Batterham PJ, Fairweather-Schmidt AK, Butterworth P, et al (2014). Temporal effects of separation on suicidal thoughts and behaviours. Soc
	Sci Med, 111: 58-63.
77552	Batty GD, Whitley E, Kivimaki M, et al (2010). Body mass index and attempted suicide: cohort study of 1,133,019 Swedish men. Am J Epidemiol, 172(8): 890-9.
86478	Baum N (2014). Professionals' double exposure in the shared traumatic reality of wartime: contributions to professional growth and stress. British Journal of Social Work, 44(8): 2113-34.
52683	Baume P (1988). Perspectives on youth suicide. The Australian Journal of Advanced Nursing, 5(3): 40-8.
52465	Bearman PS, Moody J (2004). Suicide and friendships among American adolescents. Am J Public Health, 94(1): 89-95.
55082	Bebbington PE, Cooper C, Minot S, et al (2009). Suicide attempts, gender, and sexual abuse: data from the 2000 British Psychiatric Morbidity Survey. Am J Psychiatry, 166: 1135-40.
79639	Becerra MB, Becerra BJ, Hassija CM, et al (2016). Unmet mental healthcare need and suicidal ideation among U.S. veterans. Am J Prev Med, 51(1): 90-4.
52998	Beck, AT, Steer RA, Kovacs M, et al (1985). Hopelessness and eventual suicide: a 10-year prospective study of patients hospitalized with suicidal ideation. Am J Psychiatry, 142(5): 559-63.
77553	Behera C, Krishna K, Singh HR (2013). Antitubercular drug-induced violent suicide of a hospitalised patient. BMJ. Retrieved 25 February 2016, from doi:10.1136/bcr-2012-201469
79940	Belik S-L, Stein MB, Asmundson GJG, et al (2009). Relation between traumatic events and suicide attempts in Canadian military personnel. Can J Psychiatry, 54(2): 93-104.
54291	Bell GS, Gaitatzis A, Bell CL, et al (2009). Suicide in people with epilepsy: how great is the risk? Epilepsia, 50(8): 1933-42.
53455	Bell JB, Nye EC (2007). Specific symptoms predict suicidal ideation in Vietnam combat veterans with chronic post-traumatic stress disorder. Military Medicine, 172(11): 1144-7.
53451	Benda BB (2005). Gender differences in predictors of suicidal thoughts and attempts among homeless veterans that abuse substances. Suicide & Life-Threatening Behaviour, 35(1): 106-16.
86475	Ben-Ezra M, Palgi Y, Essar N, et al (2008). Acute stress symptoms, dissociation, and depression among rescue personnel 24 hours after the Bet-Yehoshua train crash: the effects of exposure to dead bodies. Prehosp Disast Med, 23(5): 461-5.
54733	Bennett KJ, Brown KS, Boyle M, et al (2003). Does low reading achievement at school entry cause conduct problems? Soc Sci Med, 56: 2443-8.
79943	Beristianos MH, Maguen S, Neylan TC, et al (2016). Trauma exposure and risk of suicidal ideation among older adults. Am J Geriatr Psychiatry, 24(8): 639-43.
79746	Beristianos MH, Maguen S, Neylan TC, et al (2016). Trauma exposure and risk of suicidal ideation among ethnically diverse adults. Depress Anxiety, 33: 495-501.
77598	Bernert RA, Kim JS, Iwata NG, et al (2015). Sleep disturbances as an evidence-based suicide risk factor. Curr Psychiatr Rep, 17: 15.
83739	Berninger A, Webber MP, Cohen HW, et al (2010). Trends of elevated PTSD risk in firefighters exposed to the World Trade Center disaster: 2001-2005. Public Health Reports, 125(4): 556-66.
52461	Bertolote JM, Fleischmann A, De Leo D, et al (2005). Suicide attempts, plans, and ideation in culturally diverse sites: the WHO SUPRE-MISS community survey. Psychological Medicine, 35: 1457-65.

Betz ME, Valley MA, Lowen-Stein SR, et al (2011). Elevated suicide rates at high altitude: sociodemographic and health issues may be to blame. Suicide & Life-Threatening Behaviour, 41(5): 563-73.  77800 Beyer JL, Weisler RH (2016). Suicide behaviours in bipolar disorder: a review and update for the children. Psychiatr Clin N Am, 39(1): 111-23.  Bhatti JA, Nathens AB, Thiruchelvam D, et al (2016). Self-harm emergencies after bariatric surgery: a population-based cohort study. JAMA Surg. 151(3): 226-32.  Bhui K, McKenzie K, Rasul F (2007). Rates, risk factors & methods of self harm among minority ethnic groups in the UK: a systematic review. BMC Public Health, 7: 336.  Bhui K, McKenzie K, Rasul F (2007). Rates, risk factors & methods of self harm among minority ethnic groups in the UK: a systemic review. BMC Public Health, 7: 336.  Biglan K, Shoulson I (2007). [Comment] Juvenile-onset Huntington disease. A matter of perspective. Arch Neurol, 64: 783-4. Comment on ID: 54761.  Billie-Brahe U, Schmidtke A, Kerkhof AJFM, et al (1995). Background and introduction to the WHO/EURO Multicentre Study on parasuicide. Crisis, 16(2): 72-84.  Bjorkenstam C, Moller J, Ringback G, et al (2013). An association between initiation of selective serotonin reuptake inhibitors and suicide-A nationwide register-based case-crossover study. PLoS One, 8(9): e73973.  Blum K, Olive MF, Wang KK, et al (2013). Hypothesizing that designer drugs containing cathinones (*bath salts") have profound neuro-inflammatory effects and dangerous neurotoxic response following human consumption. Medical Hypotheses, 81(3): 450-5.  Bockholdt B, Klug E, Schneider V (2001). Suicide through doxylamine poisoning. Forensic Sci Int, 119: 138-40.  Boehmer TK, Flanders WD, McGeehin MA, et al (2004). Postservice mortality in Vietnam veterans: 30-year follow-up. Arch Intern Med, 164: 1908-16.  Bolton JM, Gunnell D, Turecki G (2015). Suicide risk assessment and intervention in people with mental illness. BMJ, 351: h4978.  Bolton JM, Malld R, Chaleau D, et al (2015). Ri		T
blame, Suícide & Life-Threatening Behaviour, 41(5): 563-73.  Beyer JL, Weisler RH (2016). Suicide behaviours in bipolar disorder: a review and update for the children. Psychiatr Clin N Am, 39(1): 111-23.  Bhatti JA, Nathens AB, Thiruchelvam D, et al (2016). Self-harm emergencies after bariatric surgery: a population-based cohort study. JAMA Surg. 151(3): 226-32.  Bhui K, McKenzie K, Rasul F (2007). Rates, risk factors & methods of self harm among minority ethnic groups in the UK: a systematic review. BMC Public Health, 7: 336.  Bhui K, McKenzie K, Rasul F (2007). Rates, risk factors & methods of self harm among minority ethnic groups in the UK: a systemic review. BMC Public Health, 7: 336.  Biglan K, Shoulson I (2007). [Comment] Juvenile-onset Huntington disease. A matter of perspective. Arch Neurol, 64: 783-4. Comment on ID: 54761.  Bille-Brahe U, Schmidtke A, Kerkhof AJFM, et al (1995). Background and introduction to the WHO/EURO Multicentre Study on parasulcide. Crisis, 16(2): 72-84.  Björkenstam C, Moller J, Ringback G, et al (2013). An association between initiation of selective serotonin reuptake inhibitors and suicide- A nationwide register-based case-crossover study. PLoS One, 8(9): e73973.  Blum K, Olive MF, Wang KK, et al (2013). Hypothesizing that designer drugs containing cathinones ("bath salts") have profound neuro-inflammatory effects and dangerous neurotoxic response following human consumption. Medical Hypotheses, 81(3): 450-5.  Bockholdt B, Klug E, Schneider V (2001). Suicide through doxylamine poisoning. Forensic Sci Int. 119: 138-40.  Boehmer TK, Flanders WD, McGeehin MA, et al (2004). Postservice mortality in Vietnam veterans: 30-year follow-up. Arch Intern Med, 164: 1908-16.  Bolton JM, Gunnell D, Turecki G (2015). Suicide risk assessment and intervention in people with mental illness. BMJ, 351: h4978.  Bolton JM, Walld R, Chafeau D, et al (2015). Risk factors for twelve-month suicide attempts associated with physical disorders: apopulation-based, balancing score-matched analysis. Psycholo		Betz ME, Valley MA, Lowen-Stein SR, et al (2011). Elevated suicide
President Pres	//841	
review and update for the children. Psychiatr Clin N Am. 39(1): 111-23. Bhatti JA, Nathens AB, Thiruchelvam D, et al (2016). Self-harm emergencies after bariatric surgery: a population-based cohort study. JAMA Surg. 151(3): 226-32. Bhui K, McKenzie K, Rasul F (2007). Rates, risk factors & methods of self harm among minority ethnic groups in the UK: a systematic review. BMC Public Health, 7: 336. Bhui K, McKenzie K, Rasul F (2007). Rates, risk factors & methods of self harm among minority ethnic groups in the UK: a systemic review. BMC Public Health, 7: 336. Biglan K, Shoulson I (2007). [Comment] Juvenile-onset Huntington disease. A matter of perspective. Arch Neurol, 64: 783-4. Comment on ID: 54761. Bille-Brahe U, Schmidtke A, Kerkhof AJFM, et al (1995). Background and introduction to the WHO/EURO Multicentre Study on parasuicide. Crisis, 16(2): 72-84. Bjorkenstam C, Moller J, Ringback G, et al (2013). An association between initiation of selective serotonin reuptake inhibitors and suicide-A nationwide register-based case-crossover study. PLoS One, 8(9): e73973. Blum K, Olive MF, Wang KK, et al (2013). Hypothesizing that designer drugs containing cathinones ("bath salts") have profound neuro-inflammatory effects and dangerous neurotoxic response following human consumption. Medical Hypotheses, 81(3): 450-5. Bockholdt B, Klug E, Schneider V (2001). Suicide through doxylamine poisoning. Forensic Sci Int, 119: 138-40. Boehmer TK, Flanders WD, McGeehin MA, et al (2004). Postservice mortality in Vietnam veterans: 30-year follow-up. Arch Intern Med, 164: 1908-16. Bölton JM, Pagura J, Enns MW, et al (2010). A population-based longitudinal study of risk factors for suicide attempts in major depressive disorder. J Psychiatr Res, 44(13): 817-26. Bolton JM, Walld R, Chateau D, et al (2015). Risk of suicide and suicide attempts associated with physical disorders: a population-based, balancing score-matched analysis. Psychological Med, 45(3): 495-504. Borges G, Angst J, Nock MK, et al (2006). Risk factors for twelve-month s		
emergencies after bariatric surgery: a population-based cohort study. JAMA Surg. 151(3): 226-32.  Bhui K, McKenzie K, Rasul F (2007). Rates, risk factors & methods of self harm among minority ethnic groups in the UK: a systematic review. BMC Public Health, 7: 336.  Bhui K, McKenzie K, Rasul F (2007). Rates, risk factors & methods of self harm among minority ethnic groups in the UK: a systemic review. BMC Public Health, 7: 336.  Biglan K, Shoulson I (2007). [Comment] Juvenile-onset Huntington disease. A matter of perspective. Arch Neurol, 64: 783-4. Comment on ID: 54761.  Bille-Brahe U, Schmidtke A, Kerkhof AJFM, et al (1995). Background and introduction to the WHO/EURO Multicentre Study on parasulcide. Crisis, 16(2): 72-84.  Bijorkenstam C, Moller J, Ringback G, et al (2013). An association between initiation of selective serotonin reuptake inhibitors and suicide- A nationwide register-based case-crossover study. PLoS One, 8(9): e73973.  Blum K, Olive MF, Wang KK, et al (2013). Hypothesizing that designer drugs containing cathinones ("bath salts") have profound neuro-inflammatory effects and dangerous neurotoxic response following human consumption. Medical Hypotheses, 81(3): 450-5.  Bockholdt B, Klug E, Schneider V (2001). Suicide through doxylamine poisoning. Forensic Sci Int, 119: 138-40.  Boehmer TK, Flanders WD, McGeehin MA, et al (2004). Postservice mortality in Vietnam veterans: 30-year follow-up. Arch Intern Med, 164: 1908-16.  Bolton JM, Pagura J, Enns MW, et al (2015). Suicide risk assessment and intervention in people with mental illness. BMJ, 351: h4978.  Bolton JM, Pagura J, Enns MW, et al (2015). Risk of suicide and suicide attempts associated with physical disorders: a population-based longitudinal study of risk factors for suicide attempts in major depressive disorder. J Psychiatr Res, 44(13): 817-26.  Bolton JM, Walld R, Chateau D, et al (2015). Risk of suicide and suicide attempts associated with physical disorders: a population-based, balancing score-matched analysis. Psychological Med, 45(3)	77800	review and update for the children. Psychiatr Clin N Am, 39(1): 111-23.
Bhui K, McKenzie K, Rasul F (2007). Rates, risk factors & methods of self harm among minority ethnic groups in the UK: a systematic review. BMC Public Health, 7: 336.  Bhui K, McKenzie K, Rasul F (2007). Rates, risk factors & methods of self harm among minority ethnic groups in the UK: a systemic review. BMC Public Health, 7: 336.  Biglan K, Shoulson I (2007). [Comment] Juvenile-onset Huntington disease. A matter of perspective. Arch Neurol, 64: 783-4. Comment on ID: 54761.  Bille-Brahe U, Schmidtke A, Kerkhof AJFM, et al (1995). Background and introduction to the WHO/EURO Multicentre Study on parasuicide. Crisis, 16(2): 72-84.  Bjorkenstam C, Moller J, Ringback G, et al (2013). An association between initiation of selective serotonin reuptake inhibitors and suicide-A nationwide register-based case-crossover study. PLoS One, 8(9): e73973.  Blum K, Olive MF, Wang KK, et al (2013). Hypothesizing that designer drugs containing cathinones ("bath salts") have profound neuro-inflammatory effects and dangerous neurotoxic response following human consumption. Medical Hypotheses, 81(3): 450-5.  Bockholdt B, Klug E, Schneider V (2001). Suicide through doxylamine poisoning. Forensic Sci Int, 119: 138-40.  Boehmer TK, Flanders WD, McGenhin MA, et al (2004). Postservice mortality in Vietnam veterans: 30-year follow-up. Arch Intern Med, 164: 1908-16.  Bolton JM, Gunnell D, Turecki G (2015). Suicide risk assessment and intervention in people with mental illness. BMJ, 351: h4978.  Bolton JM, Pagura J, Enns MW, et al (2010). A population-based longitudinal study of risk factors for suicide attempts in major depressive disorder. J Psychiatr Res, 44(13): 817-26.  Bolton JM, Walld R, Chateau D, et al (2015). Risk of suicide and suicide attempts in the National Comorbidity Survey Replication (NCS-R). Psychol Med, 36(12): 1747-57.  Borges G, Angts J, Nock MK, et al (2006). Risk factors for twelve-month suicide attempts in the National Comorbidity Survey Replication (NCS-R). Psychol Med, 36(12): 1747-57.  Borges G, Angts J, Nock MK,	77040	• • • • • • • • • • • • • • • • • • • •
Bhui K, McKenzie K, Rasul F (2007). Rates, risk factors & methods of self harm among minority ethnic groups in the UK: a systematic review. BMC Public Health, 7: 336.  Bhui K, McKenzie K, Rasul F (2007). Rates, risk factors & methods of self harm among minority ethnic groups in the UK: a systemic review. BMC Public Health, 7: 336.  Biglan K, Shoulson I (2007). [Comment] Juvenile-onset Huntington disease. A matter of perspective. Arch Neurol, 64: 783-4. Comment on ID: 54761.  Billie-Brahe U, Schmidtke A, Kerkhof AJFM, et al (1995). Background and introduction to the WHO/EURO Multicentre Study on parasuicide. Crisis, 16(2): 72-84.  Bjorkenstam C, Moller J, Ringback G, et al (2013). An association between initiation of selective serotonin reuptake inhibitors and suicide-A nationwide register-based case-crossover study. PLoS One, 8(9): e73973.  Blum K, Olive MF, Wang KK, et al (2013). Hypothesizing that designer drugs containing cathinones ("bath salts") have profound neuro-inflammatory effects and dangerous neurotoxic response following human consumption. Medical Hypotheses, 81(3): 450-5.  Bockholdt B, Klug E, Schneider V (2001). Suicide through doxylamine poisoning. Forensic Sci Int, 119: 138-40.  Boehmer TK, Flanders WD, McGeehin MA, et al (2004). Postservice mortality in Vietnam veterans: 30-year follow-up. Arch Intern Med, 164: 1908-16.  Bolton JM, Gunnell D, Turecki G (2015). Suicide risk assessment and intervention in people with mental illness. BMJ, 351: h4978.  Bolton JM, Pagura J, Enns MW, et al (2010). A population-based longitudinal study of risk factors for suicide attempts in major depressive disorder: J Psychiatr Res, 44(13): 817-26.  Bolton JM, Walld R, Chateau D, et al (2015). Risk of suicide and suicide attempts associated with physical disorders: a population-based, balancing score-matched analysis. Psychological Med, 45(3): 495-504.  Borges G, Angst J, Nock MK, et al (2006). Risk factors for twelve-month suicide attempts in the National Comorbidity Survey Replication (NCS-R). Psychol Med, 36(12)	77943	
self harm among minority ethnic groups in the UK: a systematic review. BMC Public Health, 7: 336.  Bhui K, McKenzie K, Rasul F (2007). Rates, risk factors & methods of self harm among minority ethnic groups in the UK: a systemic review. BMC Public Health, 7: 336.  Biglan K, Shoulson I (2007). [Comment] Juvenile-onset Huntington disease. A matter of perspective. Arch Neurol, 64: 783-4. Comment on ID: 54761.  Bille-Brahe U, Schmidtke A, Kerkhof AJFM, et al (1995). Background and introduction to the WHO/EURO Multicentre Study on parasuicide. Crisis, 16(2): 72-84.  Bjorkenstam C, Moller J, Ringback G, et al (2013). An association between initiation of selective serotonin reuptake inhibitors and suicide-A nationwide register-based case-crossover study. PLoS One, 8(9): e73973.  Blum K, Olive MF, Wang KK, et al (2013). Hypothesizing that designer drugs containing cathinones ("bath salts") have profound neuro-inflammatory effects and dangerous neurotoxic response following human consumption. Medical Hypotheses, 81(3): 450-5.  Bockholdt B, Klug E, Schneider V (2001). Suicide through doxylamine poisoning. Forensic Sci Int, 119: 138-40.  Boehmer TK, Flanders WD, McGeehin MA, et al (2004). Postservice mortality in Vietnam veterans: 30-year follow-up. Arch Intern Med, 164: 1908-16.  Bolton JM, Gunnell D, Turecki G (2015). Suicide risk assessment and intervention in people with mental illness. BMJ, 351: h4978.  Bolton JM, Pagura J, Enns MW, et al (2010). A population-based longitudinal study of risk factors for suicide attempts in major depressive disorder. J Psychiatr Res, 44(13): 817-26.  Bolton JM, Walld R, Chateau D, et al (2015). Risk of suicide and suicide attempts associated with physical disorders: a population-based longitudinal study of risk factors for suicide attempts in major depressive disorder. J Psychiatr Res, 44(13): 817-26.  Bolton JM, Walld R, Chateau D, et al (2015). Risk factors for twelve-month suicide attempts associated with physical disorders: a population-based longitudinal study of risk factors for		
Bhui K, McKenzie K, Rasul F (2007). Rates, risk factors & methods of self harm among minority ethnic groups in the UK: a systemic review. BMC Public Health, 7: 336.  Biglan K, Shoulson I (2007). [Comment] Juvenile-onset Huntington disease. A matter of perspective. Arch Neurol, 64: 783-4. Comment on ID: 54761.  Bille-Brahe U, Schmidtke A, Kerkhof AJFM, et al (1995). Background and introduction to the WHO/EURO Multicentre Study on parasuicide. Crisis, 16(2): 72-84.  Bjorkenstam C, Moller J, Ringback G, et al (2013). An association between initiation of selective serotonin reuptake inhibitors and suicide- A nationwide register-based case-crossover study. PLoS One, 8(9): e73973.  Blum K, Olive MF, Wang KK, et al (2013). Hypothesizing that designer drugs containing cathinones ("bath salts") have profound neuroinflammatory effects and dangerous neurotoxic response following human consumption. Medical Hypotheses, 81(3): 450-5.  Bockholdt B, Klug E, Schneider V (2001). Suicide through doxylamine poisoning. Forensic Sci Int, 119: 138-40.  Boehmer TK, Flanders WD, McGeehin MA, et al (2004). Postservice mortality in Vietnam veterans: 30-year follow-up. Arch Intern Med, 164: 1908-16.  Bolton JM, Gunnell D, Turecki G (2015). Suicide risk assessment and intervention in people with mental illness. BMJ, 351: h4978.  Bolton JM, Pagura J, Enns MW, et al (2010). A population-based longitudinal study of risk factors for suicide attempts in major depressive disorder: J Psychiatr Res, 44(13): 817-26.  Bolton JM, Walld R, Chateau D, et al (2015). Risk of suicide and suicide attempts associated with physical disorders: a population-based balancing score-matched analysis. Psychological Med, 45(3): 495-504.  Borges G, Angst J, Nock MK, et al (2016). Risk factors for twelve-month suicide attempts in the National Comorbidity Survey Replication (NCS-R). Psychol Med, 36(12): 1747-57.  Borges G, Anthony JC, Garrison CZ (1995). Methodological issues relevant to epidemiologic investigations of suicidal behaviors of adolescents. Epidemiologic i	52154	self harm among minority ethnic groups in the UK: a systematic review.
self harm among minority ethnic groups in the UK: a systemic review. BMC Public Health, 7: 336.  Biglan K, Shoulson I (2007). [Comment] Juvenile-onset Huntington disease. A matter of perspective. Arch Neurol, 64: 783-4. Comment on ID: 54761.  Bille-Brahe U, Schmidtke A, Kerkhof AJFM, et al (1995). Background and introduction to the WHO/EURO Multicentre Study on parasuicide. Crisis, 16(2): 72-84.  Björkenstam C, Moller J, Ringback G, et al (2013). An association between initiation of selective serotonin reuptake inhibitors and suicide- A nationwide register-based case-crossover study. PLoS One, 8(9): e73973.  Blum K, Olive MF, Wang KK, et al (2013). Hypothesizing that designer drugs containing cathinones ("bath salts") have profound neuro-inflammatory effects and dangerous neurotoxic response following human consumption. Medical Hypotheses, 81(3): 450-5.  Bockholdt B, Klug E, Schneider V (2001). Suicide through doxylamine poisoning. Forensic Sci Int., 119: 138-40.  Boehmer TK, Flanders WD, McGeehin MA, et al (2004). Postservice mortality in Vietnam veterans: 30-year follow-up. Arch Intern Med, 164: 1908-16.  Bolton JM, Gunnell D, Turecki G (2015). Suicide risk assessment and intervention in people with mental illness. BMJ, 351: h4978.  Bolton JM, Pagura J, Enns MW, et al (2010). A population-based longitudinal study of risk factors for suicide attempts in major depressive disorder. J Psychiatr Res, 44(13): 817-26.  Bolton JM, Walld R, Chateau D, et al (2015). Risk of suicide and suicide attempts associated with physical disorders: a population-based, balancing score-matched analysis. Psychological Med, 45(3): 495-504.  Borges G, Angst J, Nock MK, et al (2006). Risk factors for twelve-month suicide attempts in the National Comorbidity Survey Replication (NCS-R). Psychol Med, 36(12): 1747-57.  Borges G, Angst J, Nock MK, et al (2006). Risk factors for twelve-month suicide attempts in the National Comorbidity Survey Replication (NCS-R). Psychol Med, 36(12): 1747-57.  Borges G, Angst J, Gerrison CZ (1995). Methodolo		
Biglan K, Shoulson I (2007). [Comment] Juvenile-onset Huntington disease. A matter of perspective. Arch Neurol, 64: 783-4. Comment on ID: 54761.  Bille-Brahe U, Schmidtke A, Kerkhof AJFM, et al (1995). Background and introduction to the WHO/EURO Multicentre Study on parasuicide. Crisis, 16(2): 72-84.  Bjorkenstam C, Moller J, Ringback G, et al (2013). An association between initiation of selective serotonin reuptake inhibitors and suicide- A nationwide register-based case-crossover study. PLoS One, 8(9): e73973.  Blum K, Olive MF, Wang KK, et al (2013). Hypothesizing that designer drugs containing cathinones ("bath salts") have profound neuro-inflammatory effects and dangerous neurotoxic response following human consumption. Medical Hypotheses, 81(3): 450-5.  Bockholdt B, Klug E, Schneider V (2001). Suicide through doxylamine poisoning. Forensic Sci Int, 119: 138-40.  Boehmer TK, Flanders WD, McGeehin MA, et al (2004). Postservice mortality in Vietnam veterans: 30-year follow-up. Arch Intern Med, 164: 1908-16.  Bolton JM, Gunnell D, Turecki G (2015). Suicide risk assessment and intervention in people with mental illiness. BMJ, 351: h4978.  Bolton JM, Gunnell D, Turecki G (2015). Suicide risk assessment and intervention in people with mental illness. BMJ, 351: h4978.  Bolton JM, Walla R, Chateau D, et al (2010). A population-based longitudinal study of risk factors for suicide attempts in major depressive disorder. J Psychiatr Res, 44(13): 817-26.  Borges G, Angst J, Nock MK, et al (2006). Risk factors for twelve-month suicide attempts in the National Comorbidity Survey Replication (NCS-R). Psychol Med, 36(12): 1747-57.  Borges G, Anthony JC, Garrison CZ (1995). Methodological issues relevant to epidemiologic investigations of suicidal behaviors of adolescents. Epidemiologic Reviews, 17(1): 228-239.  Bovard RS (2008). Risk behaviors in high school and college sport. Current Sports Medicine Reports, 7(6): 359-66.  Brackbill RM, Hadler JL, Difcrande L, et al (2009). Asthma and posttraumatic stress symptoms 5 to 6	52668	self harm among minority ethnic groups in the UK: a systemic review.
disease. A matter of perspective. Arch Neurol, 64: 783-4. Comment on ID: 54761.  Bille-Brahe U, Schmidtke A, Kerkhof AJFM, et al (1995). Background and introduction to the WHO/EURO Multicentre Study on parasuicide. Crisis, 16(2): 72-84.  Bjorkenstam C, Moller J, Ringback G, et al (2013). An association between initiation of selective serotonin reuptake inhibitors and suicide- A nationwide register-based case-crossover study. PLoS One, 8(9): e73973.  Blum K, Olive MF, Wang KK, et al (2013). Hypothesizing that designer drugs containing cathinones ("bath salts") have profound neuro-inflammatory effects and dangerous neurotoxic response following human consumption. Medical Hypotheses, 81(3): 450-5.  Bockholdt B, Klug E, Schneider V (2001). Suicide through doxylamine poisoning. Forensic Sci Int, 119: 138-40.  Boehmer TK, Flanders WD, McGeehin MA, et al (2004). Postservice mortality in Vietnam veterans: 30-year follow-up. Arch Intern Med, 164: 1908-16.  Bolton JM, Gunnell D, Turecki G (2015). Suicide risk assessment and intervention in people with mental illness. BMJ, 351: h4978.  Bolton JM, Pagura J, Enns MW, et al (2010). A population-based longitudinal study of risk factors for suicide attempts in major depressive disorder. J Psychiatr Res, 44(13): 817-26.  Bolton JM, Walld R, Chateau D, et al (2015). Risk of suicide and suicide attempts associated with physical disorders: a population-based, balancing score-matched analysis. Psychological Med, 45(3): 495-504.  Borges G, Anfast J, Nock MK, et al (2006). Risk factors for twelve-month suicide attempts in the National Comorbidity Survey Replication (NCS-R). Psychol Med, 36(12): 1747-57.  Borges G, Anthony JC, Garrison CZ (1995). Methodological issues relevant to epidemiologic investigations of suicidal behaviors of adolescents. Epidemiologic Reviews, 17(1): 228-239.  Bovard RS (2008). Risk behaviors in high school and college sport. Current Sports Medicine Reports, 7(6): 359-66.  Brackbill RM, Hadler JL, DiGrande L, et al (2009). Asthma and posttraumatic stress sympt		
Bille-Brahe U, Schmidtke A, Kerkhof AJFM, et al (1995). Background and introduction to the WHO/EURO Multicentre Study on parasuicide. Crisis, 16(2): 72-84.  Björkenstam C, Moller J, Ringback G, et al (2013). An association between initiation of selective serotonin reuptake inhibitors and suicide- A nationwide register-based case-crossover study. PLoS One, 8(9): e73973.  Blum K, Olive MF, Wang KK, et al (2013). Hypothesizing that designer drugs containing cathinones ("bath salts") have profound neuro-inflammatory effects and dangerous neurotoxic response following human consumption. Medical Hypotheses, 81(3): 450-5.  Bockholdt B, Klug E, Schneider V (2001). Suicide through doxylamine poisoning. Forensic Sci Int. 119: 138-40.  Boehmer TK, Flanders WD, McGeehin MA, et al (2004). Postservice mortality in Vietnam veterans: 30-year follow-up. Arch Intern Med, 164: 1908-16.  77846  Bolton JM, Gunnell D, Turecki G (2015). Suicide risk assessment and intervention in people with mental illness. BMJ, 351: h4978.  Bolton JM, Pagura J, Enns MW, et al (2010). A population-based longitudinal study of risk factors for suicide attempts in major depressive disorder. J Psychiatr Res, 44(13): 817-26.  Bolton JM, Walld R, Chateau D, et al (2015). Risk of suicide and suicide attempts associated with physical disorders: a population-based, balancing score-matched analysis. Psychological Med, 45(3): 495-504.  Borges G, Angst J, Nock MK, et al (2006). Risk factors for twelve-month suicide attempts in the National Comorbidity Survey Replication (NCS-R). Psychol Med, 36(12): 1747-57.  Borges G, Anthony JC, Garrison CZ (1995). Methodological issues relevant to epidemiologic investigations of suicidal behaviors of adolescents. Epidemiologic Reviews, 17(1): 228-239.  Bovard RS (2008). Risk behaviors in high school and college sport. Current Sports Medicine Reports, 7(6): 359-66.  Brackbill RM, Hadler JL, DiGrande L, et al (2009). Asthma and posttraumatic stress symptoms 5 to 6 years following exposure to the World Trade Center terrorist att	54762	
Bille-Brahe U, Schmidtke A, Kerkhof AJFM, et al (1995). Background and introduction to the WHO/EURO Multicentre Study on parasuicide. Crisis, 16(2): 72-84.  Bjorkenstam C, Moller J, Ringback G, et al (2013). An association between initiation of selective serotonin reuptake inhibitors and suicide- A nationwide register-based case-crossover study. PLoS One, 8(9): e73973.  Blum K, Olive MF, Wang KK, et al (2013). Hypothesizing that designer drugs containing cathinones ("bath salts") have profound neuro-inflammatory effects and dangerous neurotoxic response following human consumption. Medical Hypotheses, 81(3): 450-5.  Bockholdt B, Klug E, Schneider V (2001). Suicide through doxylamine poisoning. Forensic Sci Int, 119: 138-40.  Boehmer TK, Flanders WD, McGeehin MA, et al (2004). Postservice mortality in Vietnam veterans: 30-year follow-up. Arch Intern Med, 164: 1908-16.  Bolton JM, Gunnell D, Turecki G (2015). Suicide risk assessment and intervention in people with mental illness. BMJ, 351: h4978.  Bolton JM, Pagura J, Enns MW, et al (2010). A population-based longitudinal study of risk factors for suicide attempts in major depressive disorder. J Psychiatr Res, 44(13): 817-26.  Bolton JM, Walld R, Chateau D, et al (2015). Risk of suicide and suicide attempts associated with physical disorders: a population-based, balancing score-matched analysis. Psychological Med, 45(3): 495-504.  Borges G, Angst J, Nock MK, et al (2006). Risk factors for twelve-month suicide attempts in the National Comorbidity Survey Replication (NCS-R). Psychol Med, 36(12): 1747-57.  Borges G, Anthony JC, Garrison CZ (1995). Methodological issues relevant to epidemiologic investigations of suicidal behaviors of adolescents. Epidemiologic Reviews, 17(1): 228-239.  Boston PF, Dursun SM, Reveley MA (1994). Other studies exist. BMJ, 309: 1228.  Bovard RS (2008). Risk behaviors in high school and college sport. Current Sports Medicine Reports, 7(6): 359-66.  Brackbill RM, Hadler JL, DiGrande L, et al (2009). Asthma and posttraumatic stress symptoms	34702	· ·
introduction to the WHO/EURO Multicentre Study on parasuicide. Crisis, 16(2): 72-84.  Bjorkenstam C, Moller J, Ringback G, et al (2013). An association between initiation of selective serotonin reuptake inhibitors and suicide- A nationwide register-based case-crossover study. PLoS One, 8(9): e73973.  Blum K, Olive MF, Wang KK, et al (2013). Hypothesizing that designer drugs containing cathinones ("bath salts") have profound neuro-inflammatory effects and dangerous neurotoxic response following human consumption. Medical Hypotheses, 81(3): 450-5.  Bockholdt B, Klug E, Schneider V (2001). Suicide through doxylamine poisoning. Forensic Sci Int, 119: 138-40.  Boehmer TK, Flanders WD, McGeehin MA, et al (2004). Postservice mortality in Vietnam veterans: 30-year follow-up. Arch Intern Med, 164: 1908-16.  Bolton JM, Gunnell D, Turecki G (2015). Suicide risk assessment and intervention in people with mental illness. BMJ, 351: h4978.  Bolton JM, Pagura J, Enns MW, et al (2010). A population-based longitudinal study of risk factors for suicide attempts in major depressive disorder. J Psychiatr Res, 44(13): 817-26.  Bolton JM, Walld R, Chateau D, et al (2015). Risk of suicide and suicide attempts associated with physical disorders: a population-based, balancing score-matched analysis. Psychological Med, 45(3): 495-504.  Borges G, Angst J, Nock MK, et al (2006). Risk factors for twelve-month suicide attempts in the National Comorbidity Survey Replication (NCS-R). Psychol Med, 36(12): 1747-57.  Borges G, Anthony JC, Garrison CZ (1995). Methodological issues relevant to epidemiologic investigations of suicidal behaviors of adolescents. Epidemiologic Reviews, 17(1): 228-239.  Bovard RS (2008). Risk behaviors in high school and college sport. Current Sports Medicine Reports, 7(6): 359-66.  Brackbill RM, Hadler JL, DiGrande L, et al (2009). Asthma and posttraumatic stress symptoms 5 to 6 years following exposure to the World Trade Center terrorist attack. JAMA, 302(5): 502-16.  Braden JB, Sullivan MD (2008). Suicidal thoughts a		
Bjorkenstam C, Moller J, Ringback G, et al (2013). An association between initiation of selective serotonin reuptake inhibitors and suicide- A nationwide register-based case-crossover study. PLoS One, 8(9): e73973.  Blum K, Olive MF, Wang KK, et al (2013). Hypothesizing that designer drugs containing cathinones ("bath salts") have profound neuro-inflammatory effects and dangerous neurotoxic response following human consumption. Medical Hypotheses, 81(3): 450-5.  54041 Bockholdt B, Klug E, Schneider V (2001). Suicide through doxylamine poisoning. Forensic Sci Int, 119: 138-40.  Boehmer TK, Flanders WD, McGeehin MA, et al (2004). Postservice mortality in Vietnam veterans: 30-year follow-up. Arch Intern Med, 164: 1908-16.  Bolton JM, Gunnell D, Turecki G (2015). Suicide risk assessment and intervention in people with mental illness. BMJ, 351: h4978.  Bolton JM, Pagura J, Enns MW, et al (2010). A population-based longitudinal study of risk factors for suicide attempts in major depressive disorder. J Psychiatr Res, 44(13): 817-26.  Bolton JM, Walld R, Chateau D, et al (2015). Risk of suicide and suicide attempts associated with physical disorders: a population-based, balancing score-matched analysis. Psychological Med, 45(3): 495-504.  Borges G, Angst J, Nock MK, et al (2006). Risk factors for twelve-month suicide attempts in the National Comorbidity Survey Replication (NCS-R). Psychol Med, 36(12): 1747-57.  Borges G, Anthony JC, Garrison CZ (1995). Methodological issues relevant to epidemiologic investigations of suicidal behaviors of adolescents. Epidemiologic Reviews, 17(1): 228-239.  Bovard RS (2008). Risk behaviors in high school and college sport. Current Sports Medicine Reports, 7(6): 359-66.  Brackbill RM, Hadler JL, DiGrande L, et al (2009). Asthma and posttraumatic stress symptoms 5 to 6 years following exposure to the World Trade Center terrorist attack. JAMA, 302(5): 502-16.  Braden JB, Sullivan MD (2008). Suicidal thoughts and behavior among adults with self-reported pain conditions in the National Como	55081	
between initiation of selective serotonin reuptake inhibitors and suicide- A nationwide register-based case-crossover study. PLoS One, 8(9): e73973.  Blum K, Olive MF, Wang KK, et al (2013). Hypothesizing that designer drugs containing cathinones ("bath salts") have profound neuro-inflammatory effects and dangerous neurotoxic response following human consumption. Medical Hypotheses, 81(3): 450-5.  Bockholdt B, Klug E, Schneider V (2001). Suicide through doxylamine poisoning. Forensic Sci Int, 119: 138-40.  Boehmer TK, Flanders WD, McGeehin MA, et al (2004). Postservice mortality in Vietnam veterans: 30-year follow-up. Arch Intern Med, 164: 1908-16.  Bolton JM, Gunnell D, Turecki G (2015). Suicide risk assessment and intervention in people with mental illness. BMJ, 351: h4978.  Bolton JM, Pagura J, Enns MW, et al (2010). A population-based longitudinal study of risk factors for suicide attempts in major depressive disorder. J Psychiatr Res, 44(13): 817-26.  Bolton JM, Walld R, Chateau D, et al (2015). Risk of suicide and suicide attempts associated with physical disorders: a population-based, balancing score-matched analysis. Psychological Med, 45(3): 495-504.  Borges G, Angst J, Nock MK, et al (2006). Risk factors for twelve-month suicide attempts in the National Comorbidity Survey Replication (NCS-R). Psychol Med, 36(12): 1747-57.  Borges G, Anthony JC, Garrison CZ (1995). Methodological issues relevant to epidemiologic investigations of suicidal behaviors of adolescents. Epidemiologic Reviews, 17(1): 228-239.  Boston PF, Dursun SM, Reveley MA (1994). Other studies exist. BMJ, 309: 1228.  Bovard RS (2008). Risk behaviors in high school and college sport. Current Sports Medicine Reports, 7(6): 359-66.  Brackbill RM, Hadler JL, DiGrande L, et al (2009). Asthma and posttraumatic stress symptoms 5 to 6 years following exposure to the World Trade Center terrorist attack. JAMA, 302(5): 502-16.  Braden JB, Sullivan MD (2008). Suicidal thoughts and behavior among adults with self-reported pain conditions in the Nation		
nationwide register-based case-crossover study. PLoS One, 8(9): e73973.  Blum K, Olive MF, Wang KK, et al (2013). Hypothesizing that designer drugs containing cathinones ("bath salts") have profound neuro-inflammatory effects and dangerous neurotoxic response following human consumption. Medical Hypotheses, 81(3): 450-5.  Bockholdt B, Klug E, Schneider V (2001). Suicide through doxylamine poisoning. Forensic Sci Int, 119: 138-40.  Boehmer TK, Flanders WD, McGeehin MA, et al (2004). Postservice mortality in Vietnam veterans: 30-year follow-up. Arch Intern Med, 164: 1908-16.  Bolton JM, Gunnell D, Turecki G (2015). Suicide risk assessment and intervention in people with mental illness. BMJ, 351: h4978.  Bolton JM, Pagura J, Enns MW, et al (2010). A population-based longitudinal study of risk factors for suicide attempts in major depressive disorder. J Psychiatr Res, 44(13): 817-26.  Bolton JM, Walld R, Chateau D, et al (2015). Risk of suicide and suicide attempts associated with physical disorders: a population-based, balancing score-matched analysis. Psychological Med, 45(3): 495-504.  Borges G, Angst J, Nock MK, et al (2006). Risk factors for twelve-month suicide attempts in the National Comorbidity Survey Replication (NCS-R). Psychol Med, 36(12): 1747-57.  Borges G, Anthony JC, Garrison CZ (1995). Methodological issues relevant to epidemiologic investigations of suicidal behaviors of adolescents. Epidemiologic Reviews, 17(1): 228-239.  Boston PF, Dursun SM, Reveley MA (1994). Other studies exist. BMJ, 309: 1228.  Bovard RS (2008). Risk behaviors in high school and college sport. Current Sports Medicine Reports, 7(6): 359-66.  Brackbill RM, Hadler JL, DiGrande L, et al (2009). Asthma and posttraumatic stress symptoms 5 to 6 years following exposure to the World Trade Center terrorist attack. JAMA, 302(5): 502-16.  Braden JB, Sullivan MD (2008). Suicidal thoughts and behavior among adults with self-reported pain conditions in the National Comorbidity		
nationwide register-based case-crossover study. PLos One, 8(9): e73973.  Blum K, Olive MF, Wang KK, et al (2013). Hypothesizing that designer drugs containing cathinones ("bath salts") have profound neuro-inflammatory effects and dangerous neurotoxic response following human consumption. Medical Hypotheses, 81(3): 450-5.  Bockholdt B, Klug E, Schneider V (2001). Suicide through doxylamine poisoning. Forensic Sci Int, 119: 138-40.  Boehmer TK, Flanders WD, McGeehin MA, et al (2004). Postservice mortality in Vietnam veterans: 30-year follow-up. Arch Intern Med, 164: 1908-16.  Bolton JM, Gunnell D, Turecki G (2015). Suicide risk assessment and intervention in people with mental illness. BMJ, 351: h4978.  Bolton JM, Pagura J, Enns MW, et al (2010). A population-based longitudinal study of risk factors for suicide attempts in major depressive disorder. J Psychiatr Res, 44(13): 817-26.  Bolton JM, Walld R, Chateau D, et al (2015). Risk of suicide and suicide attempts associated with physical disorders: a population-based, balancing score-matched analysis. Psychological Med, 45(3): 495-504.  Borges G, Angst J, Nock MK, et al (2006). Risk factors for twelve-month suicide attempts in the National Comorbidity Survey Replication (NCS-R). Psychol Med, 36(12): 1747-57.  Borges G, Anthony JC, Garrison CZ (1995). Methodological issues relevant to epidemiologic investigations of suicidal behaviors of adolescents. Epidemiologic Reviews, 17(1): 228-239.  Boston PF, Dursun SM, Reveley MA (1994). Other studies exist. BMJ, 309: 1228.  Bovard RS (2008). Risk behaviors in high school and college sport. Current Sports Medicine Reports, 7(6): 359-66.  Brackbill RM, Hadler JL, DiGrande L, et al (2009). Asthma and posttraumatic stress symptoms 5 to 6 years following exposure to the World Trade Center terrorist attack. JAMA, 302(5): 502-16.  Braden JB, Sullivan MD (2008). Suicidal thoughts and behavior among adults with self-reported pain conditions in the National Comorbidity	77370	
Blum K, Olive MF, Wang KK, et al (2013). Hypothesizing that designer drugs containing cathinones ("bath salts") have profound neuroinflammatory effects and dangerous neurotoxic response following human consumption. Medical Hypotheses, 81(3): 450-5.  Bockholdt B, Klug E, Schneider V (2001). Suicide through doxylamine poisoning. Forensic Sci Int, 119: 138-40.  Boehmer TK, Flanders WD, McGeehin MA, et al (2004). Postservice mortality in Vietnam veterans: 30-year follow-up. Arch Intern Med, 164: 1908-16.  Bolton JM, Gunnell D, Turecki G (2015). Suicide risk assessment and intervention in people with mental illness. BMJ, 351: h4978.  Bolton JM, Pagura J, Enns MW, et al (2010). A population-based longitudinal study of risk factors for suicide attempts in major depressive disorder. J Psychiatr Res, 44(13): 817-26.  Bolton JM, Walld R, Chateau D, et al (2015). Risk of suicide and suicide attempts associated with physical disorders: a population-based, balancing score-matched analysis. Psychological Med, 45(3): 495-504.  Borges G, Angst J, Nock MK, et al (2006). Risk factors for twelve-month suicide attempts in the National Comorbidity Survey Replication (NCS-R). Psychol Med, 36(12): 1747-57.  Borges G, Anthony JC, Garrison CZ (1995). Methodological issues relevant to epidemiologic investigations of suicidal behaviors of adolescents. Epidemiologic Reviews, 17(1): 228-239.  Boston PF, Dursun SM, Reveley MA (1994). Other studies exist. BMJ, 309: 1228.  Bovard RS (2008). Risk behaviors in high school and college sport. Current Sports Medicine Reports, 7(6): 359-66.  Brackbill RM, Hadler JL, DiGrande L, et al (2009). Asthma and posttraumatic stress symptoms 5 to 6 years following exposure to the World Trade Center terrorist attack. JAMA, 302(5): 502-16.  Braden JB, Sullivan MD (2008). Suicidal thoughts and behavior among adults with self-reported pain conditions in the National Comorbidity		
drugs containing cathinones ("bath salts") have profound neuro- inflammatory effects and dangerous neurotoxic response following human consumption. Medical Hypotheses, 81(3): 450-5.  Bockholdt B, Klug E, Schneider V (2001). Suicide through doxylamine poisoning. Forensic Sci Int, 119: 138-40.  Boehmer TK, Flanders WD, McGeehin MA, et al (2004). Postservice mortality in Vietnam veterans: 30-year follow-up. Arch Intern Med, 164: 1908-16.  Bolton JM, Gunnell D, Turecki G (2015). Suicide risk assessment and intervention in people with mental illness. BMJ, 351: h4978.  Bolton JM, Pagura J, Enns MW, et al (2010). A population-based longitudinal study of risk factors for suicide attempts in major depressive disorder. J Psychiatr Res, 44(13): 817-26.  Bolton JM, Walld R, Chateau D, et al (2015). Risk of suicide and suicide attempts associated with physical disorders: a population-based, balancing score-matched analysis. Psychological Med, 45(3): 495-504.  Borges G, Angst J, Nock MK, et al (2006). Risk factors for twelve-month suicide attempts in the National Comorbidity Survey Replication (NCS-R). Psychol Med, 36(12): 1747-57.  Borges G, Anthony JC, Garrison CZ (1995). Methodological issues relevant to epidemiologic investigations of suicidal behaviors of adolescents. Epidemiologic Reviews, 17(1): 228-239.  Boston FF, Dursun SM, Reveley MA (1994). Other studies exist. BMJ, 309: 1228.  Bovard RS (2008). Risk behaviors in high school and college sport. Current Sports Medicine Reports, 7(6): 359-66.  Brackbill RM, Hadler JL, DiGrande L, et al (2009). Asthma and posttraumatic stress symptoms 5 to 6 years following exposure to the World Trade Center terrorist attack. JAMA, 302(5): 502-16.  Braden JB, Sullivan MD (2008). Suicidal thoughts and behavior among adults with self-reported pain conditions in the National Comorbidity		
inflammatory effects and dangerous neurotoxic response following human consumption. Medical Hypotheses, 81(3): 450-5.  Bockholdt B, Klug E, Schneider V (2001). Suicide through doxylamine poisoning. Forensic Sci Int, 119: 138-40.  Boehmer TK, Flanders WD, McGeehin MA, et al (2004). Postservice mortality in Vietnam veterans: 30-year follow-up. Arch Intern Med, 164: 1908-16.  Bolton JM, Gunnell D, Turecki G (2015). Suicide risk assessment and intervention in people with mental illness. BMJ, 351: h4978.  Bolton JM, Pagura J, Enns MW, et al (2010). A population-based longitudinal study of risk factors for suicide attempts in major depressive disorder. J Psychiatr Res, 44(13): 817-26.  Bolton JM, Walld R, Chateau D, et al (2015). Risk of suicide and suicide attempts associated with physical disorders: a population-based, balancing score-matched analysis. Psychological Med, 45(3): 495-504.  Borges G, Angst J, Nock MK, et al (2006). Risk factors for twelve-month suicide attempts in the National Comorbidity Survey Replication (NCS-R). Psychol Med, 36(12): 1747-57.  Borges G, Anthony JC, Garrison CZ (1995). Methodological issues relevant to epidemiologic investigations of suicidal behaviors of adolescents. Epidemiologic Investigations of suicidal behaviors of adolescents. Epidemiologic Reviews, 17(1): 228-239.  Boston PF, Dursun SM, Reveley MA (1994). Other studies exist. BMJ, 309: 1228.  Bovard RS (2008). Risk behaviors in high school and college sport. Current Sports Medicine Reports, 7(6): 359-66.  Brackbill RM, Hadler JL, DiGrande L, et al (2009). Asthma and posttraumatic stress symptoms 5 to 6 years following exposure to the World Trade Center terrorist attack. JAMA, 302(5): 502-16.  Braden JB, Sullivan MD (2008). Suicidal thoughts and behavior among adults with self-reported pain conditions in the National Comorbidity		drugs containing cathingnes ("hath salts") have profound neuro-
human consumption. Medical Hypotheses, 81(3): 450-5.  Bockholdt B, Klug E, Schneider V (2001). Suicide through doxylamine poisoning. Forensic Sci Int, 119: 138-40.  Boehmer TK, Flanders WD, McGeehin MA, et al (2004). Postservice mortality in Vietnam veterans: 30-year follow-up. Arch Intern Med, 164: 1908-16.  T7846  Bolton JM, Gunnell D, Turecki G (2015). Suicide risk assessment and intervention in people with mental illness. BMJ, 351: h4978.  Bolton JM, Pagura J, Enns MW, et al (2010). A population-based longitudinal study of risk factors for suicide attempts in major depressive disorder. J Psychiatr Res, 44(13): 817-26.  Bolton JM, Walld R, Chateau D, et al (2015). Risk of suicide and suicide attempts associated with physical disorders: a population-based, balancing score-matched analysis. Psychological Med, 45(3): 495-504.  Borges G, Angst J, Nock MK, et al (2006). Risk factors for twelve-month suicide attempts in the National Comorbidity Survey Replication (NCS-R). Psychol Med, 36(12): 1747-57.  Borges G, Anthony JC, Garrison CZ (1995). Methodological issues relevant to epidemiologic investigations of suicidal behaviors of adolescents. Epidemiologic Reviews, 17(1): 228-239.  Boston PF, Dursun SM, Reveley MA (1994). Other studies exist. BMJ, 309: 1228.  Bovard RS (2008). Risk behaviors in high school and college sport. Current Sports Medicine Reports, 7(6): 359-66.  Brackbill RM, Hadler JL, DiGrande L, et al (2009). Asthma and posttraumatic stress symptoms 5 to 6 years following exposure to the World Trade Center terrorist attack. JAMA, 302(5): 502-16.  Braden JB, Sullivan MD (2008). Suicidal thoughts and behavior among adults with self-reported pain conditions in the National Comorbidity	78099	
Bockholdt B, Klug E, Schneider V (2001). Suicide through doxylamine poisoning. Forensic Sci Int, 119: 138-40.  Boehmer TK, Flanders WD, McGeehin MA, et al (2004). Postservice mortality in Vietnam veterans: 30-year follow-up. Arch Intern Med, 164: 1908-16.  77846  Bolton JM, Gunnell D, Turecki G (2015). Suicide risk assessment and intervention in people with mental illness. BMJ, 351: h4978.  Bolton JM, Pagura J, Enns MW, et al (2010). A population-based longitudinal study of risk factors for suicide attempts in major depressive disorder. J Psychiatr Res, 44(13): 817-26.  Bolton JM, Walld R, Chateau D, et al (2015). Risk of suicide and suicide attempts associated with physical disorders: a population-based, balancing score-matched analysis. Psychological Med, 45(3): 495-504.  Borges G, Angst J, Nock MK, et al (2006). Risk factors for twelve-month suicide attempts in the National Comorbidity Survey Replication (NCS-R). Psychol Med, 36(12): 1747-57.  Borges G, Anthony JC, Garrison CZ (1995). Methodological issues relevant to epidemiologic investigations of suicidal behaviors of adolescents. Epidemiologic Reviews, 17(1): 228-239.  Boston PF, Dursun SM, Reveley MA (1994). Other studies exist. BMJ, 309: 1228.  Bovard RS (2008). Risk behaviors in high school and college sport. Current Sports Medicine Reports, 7(6): 359-66.  Brackbill RM, Hadler JL, DiGrande L, et al (2009). Asthma and posttraumatic stress symptoms 5 to 6 years following exposure to the World Trade Center terrorist attack. JAMA, 302(5): 502-16.  Braden JB, Sullivan MD (2008). Suicidal thoughts and behavior among adults with self-reported pain conditions in the National Comorbidity		•
Boehmer TK, Flanders WD, McGeehin MA, et al (2004). Postservice mortality in Vietnam veterans: 30-year follow-up. Arch Intern Med, 164: 1908-16.  77846 Bolton JM, Gunnell D, Turecki G (2015). Suicide risk assessment and intervention in people with mental illness. BMJ, 351: h4978.  Bolton JM, Pagura J, Enns MW, et al (2010). A population-based longitudinal study of risk factors for suicide attempts in major depressive disorder. J Psychiatr Res, 44(13): 817-26.  Bolton JM, Walld R, Chateau D, et al (2015). Risk of suicide and suicide attempts associated with physical disorders: a population-based, balancing score-matched analysis. Psychological Med, 45(3): 495-504.  Borges G, Angst J, Nock MK, et al (2006). Risk factors for twelve-month suicide attempts in the National Comorbidity Survey Replication (NCS-R). Psychol Med, 36(12): 1747-57.  Borges G, Anthony JC, Garrison CZ (1995). Methodological issues relevant to epidemiologic investigations of suicidal behaviors of adolescents. Epidemiologic Reviews, 17(1): 228-239.  Boston PF, Dursun SM, Reveley MA (1994). Other studies exist. BMJ, 309: 1228.  Bovard RS (2008). Risk behaviors in high school and college sport. Current Sports Medicine Reports, 7(6): 359-66.  Brackbill RM, Hadler JL, DiGrande L, et al (2009). Asthma and posttraumatic stress symptoms 5 to 6 years following exposure to the World Trade Center terrorist attack. JAMA, 302(5): 502-16.  Braden JB, Sullivan MD (2008). Suicidal thoughts and behavior among adults with self-reported pain conditions in the National Comorbidity	54041	
mortality in Vietnam veterans: 30-year follow-up. Arch Intern Med, 164: 1908-16.  77846  Bolton JM, Gunnell D, Turecki G (2015). Suicide risk assessment and intervention in people with mental illness. BMJ, 351: h4978.  Bolton JM, Pagura J, Enns MW, et al (2010). A population-based longitudinal study of risk factors for suicide attempts in major depressive disorder. J Psychiatr Res, 44(13): 817-26.  Bolton JM, Walld R, Chateau D, et al (2015). Risk of suicide and suicide attempts associated with physical disorders: a population-based, balancing score-matched analysis. Psychological Med, 45(3): 495-504.  Borges G, Angst J, Nock MK, et al (2006). Risk factors for twelve-month suicide attempts in the National Comorbidity Survey Replication (NCS-R). Psychol Med, 36(12): 1747-57.  Borges G, Anthony JC, Garrison CZ (1995). Methodological issues relevant to epidemiologic investigations of suicidal behaviors of adolescents. Epidemiologic Reviews, 17(1): 228-239.  Boston PF, Dursun SM, Reveley MA (1994). Other studies exist. BMJ, 309: 1228.  Bovard RS (2008). Risk behaviors in high school and college sport. Current Sports Medicine Reports, 7(6): 359-66.  Brackbill RM, Hadler JL, DiGrande L, et al (2009). Asthma and posttraumatic stress symptoms 5 to 6 years following exposure to the World Trade Center terrorist attack. JAMA, 302(5): 502-16.  Braden JB, Sullivan MD (2008). Suicidal thoughts and behavior among adults with self-reported pain conditions in the National Comorbidity	34041	
1908-16.  1908-16.  Bolton JM, Gunnell D, Turecki G (2015). Suicide risk assessment and intervention in people with mental illness. BMJ, 351: h4978.  Bolton JM, Pagura J, Enns MW, et al (2010). A population-based longitudinal study of risk factors for suicide attempts in major depressive disorder. J Psychiatr Res, 44(13): 817-26.  Bolton JM, Walld R, Chateau D, et al (2015). Risk of suicide and suicide attempts associated with physical disorders: a population-based, balancing score-matched analysis. Psychological Med, 45(3): 495-504.  Borges G, Angst J, Nock MK, et al (2006). Risk factors for twelve-month suicide attempts in the National Comorbidity Survey Replication (NCS-R). Psychol Med, 36(12): 1747-57.  Borges G, Anthony JC, Garrison CZ (1995). Methodological issues relevant to epidemiologic investigations of suicidal behaviors of adolescents. Epidemiologic Reviews, 17(1): 228-239.  Boston PF, Dursun SM, Reveley MA (1994). Other studies exist. BMJ, 309: 1228.  Bovard RS (2008). Risk behaviors in high school and college sport. Current Sports Medicine Reports, 7(6): 359-66.  Brackbill RM, Hadler JL, DiGrande L, et al (2009). Asthma and posttraumatic stress symptoms 5 to 6 years following exposure to the World Trade Center terrorist attack. JAMA, 302(5): 502-16.  Braden JB, Sullivan MD (2008). Suicidal thoughts and behavior among adults with self-reported pain conditions in the National Comorbidity	40.40=	
Bolton JM, Gunnell D, Turecki G (2015). Suicide risk assessment and intervention in people with mental illness. BMJ, 351: h4978.  Bolton JM, Pagura J, Enns MW, et al (2010). A population-based longitudinal study of risk factors for suicide attempts in major depressive disorder. J Psychiatr Res, 44(13): 817-26.  Bolton JM, Walld R, Chateau D, et al (2015). Risk of suicide and suicide attempts associated with physical disorders: a population-based, balancing score-matched analysis. Psychological Med, 45(3): 495-504.  Borges G, Angst J, Nock MK, et al (2006). Risk factors for twelve-month suicide attempts in the National Comorbidity Survey Replication (NCS-R). Psychol Med, 36(12): 1747-57.  Borges G, Anthony JC, Garrison CZ (1995). Methodological issues relevant to epidemiologic investigations of suicidal behaviors of adolescents. Epidemiologic Reviews, 17(1): 228-239.  Boston PF, Dursun SM, Reveley MA (1994). Other studies exist. BMJ, 309: 1228.  Bovard RS (2008). Risk behaviors in high school and college sport. Current Sports Medicine Reports, 7(6): 359-66.  Brackbill RM, Hadler JL, DiGrande L, et al (2009). Asthma and posttraumatic stress symptoms 5 to 6 years following exposure to the World Trade Center terrorist attack. JAMA, 302(5): 502-16.  Braden JB, Sullivan MD (2008). Suicidal thoughts and behavior among adults with self-reported pain conditions in the National Comorbidity	49495	
intervention in people with mental illness. BMJ, 351: h4978.  Bolton JM, Pagura J, Enns MW, et al (2010). A population-based longitudinal study of risk factors for suicide attempts in major depressive disorder. J Psychiatr Res, 44(13): 817-26.  Bolton JM, Walld R, Chateau D, et al (2015). Risk of suicide and suicide attempts associated with physical disorders: a population-based, balancing score-matched analysis. Psychological Med, 45(3): 495-504.  Borges G, Angst J, Nock MK, et al (2006). Risk factors for twelve-month suicide attempts in the National Comorbidity Survey Replication (NCS-R). Psychol Med, 36(12): 1747-57.  Borges G, Anthony JC, Garrison CZ (1995). Methodological issues relevant to epidemiologic investigations of suicidal behaviors of adolescents. Epidemiologic Reviews, 17(1): 228-239.  Boston PF, Dursun SM, Reveley MA (1994). Other studies exist. BMJ, 309: 1228.  Bovard RS (2008). Risk behaviors in high school and college sport. Current Sports Medicine Reports, 7(6): 359-66.  Brackbill RM, Hadler JL, DiGrande L, et al (2009). Asthma and posttraumatic stress symptoms 5 to 6 years following exposure to the World Trade Center terrorist attack. JAMA, 302(5): 502-16.  Braden JB, Sullivan MD (2008). Suicidal thoughts and behavior among adults with self-reported pain conditions in the National Comorbidity		
Bolton JM, Pagura J, Enns MW, et al (2010). A population-based longitudinal study of risk factors for suicide attempts in major depressive disorder. J Psychiatr Res, 44(13): 817-26.  Bolton JM, Walld R, Chateau D, et al (2015). Risk of suicide and suicide attempts associated with physical disorders: a population-based, balancing score-matched analysis. Psychological Med, 45(3): 495-504.  Borges G, Angst J, Nock MK, et al (2006). Risk factors for twelve-month suicide attempts in the National Comorbidity Survey Replication (NCS-R). Psychol Med, 36(12): 1747-57.  Borges G, Anthony JC, Garrison CZ (1995). Methodological issues relevant to epidemiologic investigations of suicidal behaviors of adolescents. Epidemiologic Reviews, 17(1): 228-239.  Boston PF, Dursun SM, Reveley MA (1994). Other studies exist. BMJ, 309: 1228.  Bovard RS (2008). Risk behaviors in high school and college sport. Current Sports Medicine Reports, 7(6): 359-66.  Brackbill RM, Hadler JL, DiGrande L, et al (2009). Asthma and posttraumatic stress symptoms 5 to 6 years following exposure to the World Trade Center terrorist attack. JAMA, 302(5): 502-16.  Braden JB, Sullivan MD (2008). Suicidal thoughts and behavior among adults with self-reported pain conditions in the National Comorbidity	77846	
longitudinal study of risk factors for suicide attempts in major depressive disorder. J Psychiatr Res, 44(13): 817-26.  Bolton JM, Walld R, Chateau D, et al (2015). Risk of suicide and suicide attempts associated with physical disorders: a population-based, balancing score-matched analysis. Psychological Med, 45(3): 495-504.  Borges G, Angst J, Nock MK, et al (2006). Risk factors for twelve-month suicide attempts in the National Comorbidity Survey Replication (NCS-R). Psychol Med, 36(12): 1747-57.  Borges G, Anthony JC, Garrison CZ (1995). Methodological issues relevant to epidemiologic investigations of suicidal behaviors of adolescents. Epidemiologic Reviews, 17(1): 228-239.  Boston PF, Dursun SM, Reveley MA (1994). Other studies exist. BMJ, 309: 1228.  Bovard RS (2008). Risk behaviors in high school and college sport. Current Sports Medicine Reports, 7(6): 359-66.  Brackbill RM, Hadler JL, DiGrande L, et al (2009). Asthma and posttraumatic stress symptoms 5 to 6 years following exposure to the World Trade Center terrorist attack. JAMA, 302(5): 502-16.  Braden JB, Sullivan MD (2008). Suicidal thoughts and behavior among adults with self-reported pain conditions in the National Comorbidity		
disorder. J Psychiatr Res, 44(13): 817-26.  Bolton JM, Walld R, Chateau D, et al (2015). Risk of suicide and suicide attempts associated with physical disorders: a population-based, balancing score-matched analysis. Psychological Med, 45(3): 495-504.  Borges G, Angst J, Nock MK, et al (2006). Risk factors for twelve-month suicide attempts in the National Comorbidity Survey Replication (NCS-R). Psychol Med, 36(12): 1747-57.  Borges G, Anthony JC, Garrison CZ (1995). Methodological issues relevant to epidemiologic investigations of suicidal behaviors of adolescents. Epidemiologic Reviews, 17(1): 228-239.  Boston PF, Dursun SM, Reveley MA (1994). Other studies exist. BMJ, 309: 1228.  Bovard RS (2008). Risk behaviors in high school and college sport. Current Sports Medicine Reports, 7(6): 359-66.  Brackbill RM, Hadler JL, DiGrande L, et al (2009). Asthma and posttraumatic stress symptoms 5 to 6 years following exposure to the World Trade Center terrorist attack. JAMA, 302(5): 502-16.  Braden JB, Sullivan MD (2008). Suicidal thoughts and behavior among adults with self-reported pain conditions in the National Comorbidity	77765	
attempts associated with physical disorders: a population-based, balancing score-matched analysis. Psychological Med, 45(3): 495-504.  Borges G, Angst J, Nock MK, et al (2006). Risk factors for twelve-month suicide attempts in the National Comorbidity Survey Replication (NCS-R). Psychol Med, 36(12): 1747-57.  Borges G, Anthony JC, Garrison CZ (1995). Methodological issues relevant to epidemiologic investigations of suicidal behaviors of adolescents. Epidemiologic Reviews, 17(1): 228-239.  Boston PF, Dursun SM, Reveley MA (1994). Other studies exist. BMJ, 309: 1228.  Bovard RS (2008). Risk behaviors in high school and college sport. Current Sports Medicine Reports, 7(6): 359-66.  Brackbill RM, Hadler JL, DiGrande L, et al (2009). Asthma and posttraumatic stress symptoms 5 to 6 years following exposure to the World Trade Center terrorist attack. JAMA, 302(5): 502-16.  Braden JB, Sullivan MD (2008). Suicidal thoughts and behavior among adults with self-reported pain conditions in the National Comorbidity		
balancing score-matched analysis. Psychological Med, 45(3): 495-504.  Borges G, Angst J, Nock MK, et al (2006). Risk factors for twelve-month suicide attempts in the National Comorbidity Survey Replication (NCS-R). Psychol Med, 36(12): 1747-57.  Borges G, Anthony JC, Garrison CZ (1995). Methodological issues relevant to epidemiologic investigations of suicidal behaviors of adolescents. Epidemiologic Reviews, 17(1): 228-239.  Boston PF, Dursun SM, Reveley MA (1994). Other studies exist. BMJ, 309: 1228.  Bovard RS (2008). Risk behaviors in high school and college sport. Current Sports Medicine Reports, 7(6): 359-66.  Brackbill RM, Hadler JL, DiGrande L, et al (2009). Asthma and posttraumatic stress symptoms 5 to 6 years following exposure to the World Trade Center terrorist attack. JAMA, 302(5): 502-16.  Braden JB, Sullivan MD (2008). Suicidal thoughts and behavior among adults with self-reported pain conditions in the National Comorbidity		
Borges G, Angst J, Nock MK, et al (2006). Risk factors for twelve-month suicide attempts in the National Comorbidity Survey Replication (NCS-R). Psychol Med, 36(12): 1747-57.  Borges G, Anthony JC, Garrison CZ (1995). Methodological issues relevant to epidemiologic investigations of suicidal behaviors of adolescents. Epidemiologic Reviews, 17(1): 228-239.  Boston PF, Dursun SM, Reveley MA (1994). Other studies exist. BMJ, 309: 1228.  Bovard RS (2008). Risk behaviors in high school and college sport. Current Sports Medicine Reports, 7(6): 359-66.  Brackbill RM, Hadler JL, DiGrande L, et al (2009). Asthma and posttraumatic stress symptoms 5 to 6 years following exposure to the World Trade Center terrorist attack. JAMA, 302(5): 502-16.  Braden JB, Sullivan MD (2008). Suicidal thoughts and behavior among adults with self-reported pain conditions in the National Comorbidity	77722	
<ul> <li>suicide attempts in the National Comorbidity Survey Replication (NCS-R). Psychol Med, 36(12): 1747-57.</li> <li>Borges G, Anthony JC, Garrison CZ (1995). Methodological issues relevant to epidemiologic investigations of suicidal behaviors of adolescents. Epidemiologic Reviews, 17(1): 228-239.</li> <li>Boston PF, Dursun SM, Reveley MA (1994). Other studies exist. BMJ, 309: 1228.</li> <li>Bovard RS (2008). Risk behaviors in high school and college sport. Current Sports Medicine Reports, 7(6): 359-66.</li> <li>Brackbill RM, Hadler JL, DiGrande L, et al (2009). Asthma and posttraumatic stress symptoms 5 to 6 years following exposure to the World Trade Center terrorist attack. JAMA, 302(5): 502-16.</li> <li>Braden JB, Sullivan MD (2008). Suicidal thoughts and behavior among adults with self-reported pain conditions in the National Comorbidity</li> </ul>		
Psychol Med, 36(12): 1747-57.  Borges G, Anthony JC, Garrison CZ (1995). Methodological issues relevant to epidemiologic investigations of suicidal behaviors of adolescents. Epidemiologic Reviews, 17(1): 228-239.  Boston PF, Dursun SM, Reveley MA (1994). Other studies exist. BMJ, 309: 1228.  Bovard RS (2008). Risk behaviors in high school and college sport. Current Sports Medicine Reports, 7(6): 359-66.  Brackbill RM, Hadler JL, DiGrande L, et al (2009). Asthma and posttraumatic stress symptoms 5 to 6 years following exposure to the World Trade Center terrorist attack. JAMA, 302(5): 502-16.  Braden JB, Sullivan MD (2008). Suicidal thoughts and behavior among adults with self-reported pain conditions in the National Comorbidity	40006	
Borges G, Anthony JC, Garrison CZ (1995). Methodological issues relevant to epidemiologic investigations of suicidal behaviors of adolescents. Epidemiologic Reviews, 17(1): 228-239.  Boston PF, Dursun SM, Reveley MA (1994). Other studies exist. BMJ, 309: 1228.  Bovard RS (2008). Risk behaviors in high school and college sport. Current Sports Medicine Reports, 7(6): 359-66.  Brackbill RM, Hadler JL, DiGrande L, et al (2009). Asthma and posttraumatic stress symptoms 5 to 6 years following exposure to the World Trade Center terrorist attack. JAMA, 302(5): 502-16.  Braden JB, Sullivan MD (2008). Suicidal thoughts and behavior among adults with self-reported pain conditions in the National Comorbidity	49900	• • • • • • • • • • • • • • • • • • • •
relevant to epidemiologic investigations of suicidal behaviors of adolescents. Epidemiologic Reviews, 17(1): 228-239.  Boston PF, Dursun SM, Reveley MA (1994). Other studies exist. BMJ, 309: 1228.  Bovard RS (2008). Risk behaviors in high school and college sport. Current Sports Medicine Reports, 7(6): 359-66.  Brackbill RM, Hadler JL, DiGrande L, et al (2009). Asthma and posttraumatic stress symptoms 5 to 6 years following exposure to the World Trade Center terrorist attack. JAMA, 302(5): 502-16.  Braden JB, Sullivan MD (2008). Suicidal thoughts and behavior among adults with self-reported pain conditions in the National Comorbidity		
adolescents. Epidemiologic Reviews, 17(1): 228-239.  Boston PF, Dursun SM, Reveley MA (1994). Other studies exist. BMJ, 309: 1228.  Bovard RS (2008). Risk behaviors in high school and college sport. Current Sports Medicine Reports, 7(6): 359-66.  Brackbill RM, Hadler JL, DiGrande L, et al (2009). Asthma and posttraumatic stress symptoms 5 to 6 years following exposure to the World Trade Center terrorist attack. JAMA, 302(5): 502-16.  Braden JB, Sullivan MD (2008). Suicidal thoughts and behavior among adults with self-reported pain conditions in the National Comorbidity	12291	
Boston PF, Dursun SM, Reveley MA (1994). Other studies exist. BMJ, 309: 1228.  Bovard RS (2008). Risk behaviors in high school and college sport. Current Sports Medicine Reports, 7(6): 359-66.  Brackbill RM, Hadler JL, DiGrande L, et al (2009). Asthma and posttraumatic stress symptoms 5 to 6 years following exposure to the World Trade Center terrorist attack. JAMA, 302(5): 502-16.  Braden JB, Sullivan MD (2008). Suicidal thoughts and behavior among adults with self-reported pain conditions in the National Comorbidity		
Bovard RS (2008). Risk behaviors in high school and college sport.  Current Sports Medicine Reports, 7(6): 359-66.  Brackbill RM, Hadler JL, DiGrande L, et al (2009). Asthma and posttraumatic stress symptoms 5 to 6 years following exposure to the World Trade Center terrorist attack. JAMA, 302(5): 502-16.  Braden JB, Sullivan MD (2008). Suicidal thoughts and behavior among adults with self-reported pain conditions in the National Comorbidity	1102	Boston PF, Dursun SM, Reveley MA (1994). Other studies exist. BMJ,
Current Sports Medicine Reports, 7(6): 359-66.  Brackbill RM, Hadler JL, DiGrande L, et al (2009). Asthma and posttraumatic stress symptoms 5 to 6 years following exposure to the World Trade Center terrorist attack. JAMA, 302(5): 502-16.  Braden JB, Sullivan MD (2008). Suicidal thoughts and behavior among adults with self-reported pain conditions in the National Comorbidity	7132	
Brackbill RM, Hadler JL, DiGrande L, et al (2009). Asthma and posttraumatic stress symptoms 5 to 6 years following exposure to the World Trade Center terrorist attack. JAMA, 302(5): 502-16.  Braden JB, Sullivan MD (2008). Suicidal thoughts and behavior among adults with self-reported pain conditions in the National Comorbidity	54022	
posttraumatic stress symptoms 5 to 6 years following exposure to the World Trade Center terrorist attack. JAMA, 302(5): 502-16.  Braden JB, Sullivan MD (2008). Suicidal thoughts and behavior among adults with self-reported pain conditions in the National Comorbidity		
World Trade Center terrorist attack. JAMA, 302(5): 502-16.  Braden JB, Sullivan MD (2008). Suicidal thoughts and behavior among adults with self-reported pain conditions in the National Comorbidity	63980	
Braden JB, Sullivan MD (2008). Suicidal thoughts and behavior among adults with self-reported pain conditions in the National Comorbidity		
54011 adults with self-reported pain conditions in the National Comorbidity		
· · · · · · · · · · · · · · · · · · ·	54011	, , ,

79747	Brancu M, Mann-Wrobel M, Beckham JC, et al (2016). Subthreshold posttraumatic stress disorder: a meta-analytic review of DSM-5 approach
	to measurement. Psychol Trauma, 8(2): 222-32.
54826	Breed W (1972). Five components of a basic suicide syndrome. Life-
	Threatening Behavior, 2(1): 3-18.
	Bremner JD, Shearer K, McCaffery, et al (2012). Retinoic Acid and
78064	affective disorders: the evidence for an association. J Clin Psychiatry,
	73(1): 37-50.
	Breslau N (1992). Migraine, suicidal ideation, and suicide attempts.
1741	Neurology, 42(2): 392-5.
	Breslau N, Davis GC, Andreski P (1991). Migraine, psychiatric disorders,
1742	and suicide attempts: an epidemiological study of young adults.
1742	
	Psychiatry Res, 37(1): 11-23.
45505	Brezo J, Paris J, Turecki G (2006). Personality traits as correlates of
45565	suicidal ideation, suicide attempts, and suicide completions: a systematic
	review. Acta Psychiatr Scand, 113: 180-206.
	Bronisch T, Brunner J, Bondy B, et al (2005). A multicenter study about
52680	neurobiology of suicidal behavior: design, development, and preliminary
	results. Arch Suicide Res, 9: 19-26.
	Bronisch T, Wittchen HU (1994). Suicidal ideation and suicide attempts:
4210	Comorbidity with depression, anxiety disorders, and substance abuse
	disorder. Eur Arch Psychiatry Clin Neurosci, 244: 93-98.
	Brooks SK, Dunn R, Amlot R, et al (2016). Social and occupational
84406	factors associated with psychological distress and disorder among
	disaster responders: a systematic review. BMC Psychology, 4: 18.
	Brown ES (2012). [Comment] An epidemiological approach to "steroid
77576	psychosis". Am J Psychiatry, 169(5): 447-9.
	Brown GK, Beck AT, Steer RA, et al (2000). Risk factors for suicide in
79753	psychiatric outpatients 1 20-year prospective study. J Consult Clin
10.00	Psychol, 68(3): 371-7.
	Browning SR, Westneat SC, McKnight RH (2008). Suicides among
53994	farmers in three southeaster states, 1990-1998. Journal of Agricultural
00001	Safety and Health, 14(4): 461-72.
	Brueckner RP, Lasseter KC, Lin ET, et al (1998). First-time-in humans
16206	safety and pharmacokinetics of WR238605, a new antimalarial. Am J
10200	Trop Med Hyg, 58(5): 645-49.
	Brunner J, Parhofer KG, Schwandt P, et al (2002). Cholesterol, essential
54455	fatty acids, and suicide. Pharmacopsychiatry, 35: 1-5.
77024	Bruschi Kelles SM, Diniz MF, Machado CJ, et al (2014). Mortality rate
77931	after open roux-in-Y gastric bypass: a 10-year follow-up. Braz J Med Biol
	Res, 47(7): 617-25.
70050	Bryan CJ, Cukrowicz KC (2011). Associations between types of combat
79952	violence and the acquired capability for suicide. Suicide & Life-
	Threatening Behaviour, 42(2): 126-36.
	Bryan CJ, Griffith JE, Pace BT, et al (2015). Combat exposure and risk
80090	for suicidal thoughts and behaviors among military personnel and
	veterans: a systematic review and meta-analysis. Suicide & Life-
	Threatening Behaviour, 45(5): 633-49.
81278	Bullman T, Schneiderman A, Bossarte R (2017). Suicide risk by unit
	component among veterans who served in Iraq or Afghanistan. Arch
	Suicide Res, 10: 1-10.
	Bullman TA, Kang HE (1994). Posttraumatic stress disorder and the risk
4172	of traumatic deaths among Vietnam veterans. The Journal of Nervous
	and Mental Disease, 182(11): 604-10.
0044	Bullman TA, Kang HK (1996). The risk of suicide among wounded
9341	Vietnam veterans. American Journal of Public Health, 86(5): 662-7.
L	1

55080	Burless C, De Leo D (2001). Methodological issues in community
	surveys of suicide ideators and attempters. Crisis, 22(3): 108-24.
53678	Bursztein C, Apter A (2008). Adolescent suicide. Curr Opin Psychiatry, 22: 1-6.
	Butler T, Allnutt S, Kariminia A, (2007). Mental health status of Aboriginal
52454	and non-Aboriginal Australian prisoners. Aust N Z J Psychiatry, 41: 429-
J24J4	35.
	Caldwell CB, Gottesman II (1990). Schizophrenics kill themselves too: a
1743	review of risk factors for suicide. Schizophr Bull, 16(4): 571-89.
	Callreus T, Andersen UA, Hallas J, et al (2007). Cardiovascular drugs
54034	and the risk of suicide: a nested case-control study. Eur J Clin
0.00.	Pharmacol, 63: 591-6.
50075	Canetto SS (2008). Women and suicide behavior: A cultural analysis. Am
52675	J Orthopsychiatry, 78(2): 259-66.
	Cannon J, Fitzgerald B, Seed M, et al (2015). Occupational asthma from
78012	tafenoquine in the pharmaceutical industry: implications for QSAR.
	Occup Med, 65(3): 256-8.
	Carballo JJ, Akamnonu CP, Oquendo MA (2008). Neurobiology of
52999	suicidal behavior. An integration of biological and clinical findings. Arch
	Suicide Res, 12(2): 93-110.
77795	Carlsten A, Waern M (2009). Are sedatives and hypnotics associated
77750	with increased suicide risk of suicide in the elderly? BMC Geriatr, 9: 20.
	Carpenter DJ, Fong R, Kraus JE, et al (2011). Meta-analysis of efficacy
77569	and treatment-emergent suicidality in adults by psychiatric indication and
7.7000	age subgroup following initiation of paroxetine therapy: a complete set of
	randomized placebo-controlled trails. J Clin Psychiatry, 72(11): 1503-14.
4750	Carpenter WT, Buchanan RW (1995). Schizophrenia: clinical features. HI
1758	Kaplan and BJ Sadock (Eds). Comprehensive Textbook of Psychiatry,
	6th Edition, 1 14: 889-942. Williams and Wilkins, London.  Carroll R, Metcalfe C, Gunnell D (2014). Hospital presenting self-harm
77847	and risk of fatal and non-fatal repetition: systematic review and meta-
77047	analysis. PLoS One, 9(2): e89944.
	Casey P R (1989). Personality disorder and suicide intent. Acta Psychiatr
4175	Scand, 79: 290-5.
	Cato JE, Canetto SS (2003). Attitudes and beliefs about suicidal behavior
54678	when coming out is the precipitant of the suicidal behavior. Sex Roles,
	49(9/10): 497-505.
	Cato JÉ, Canetto SS (2003). Young adults' reactions to gay and lesbian
54677	peers who became suicidal following "coming out" to their parents.
	Suicide & Life-Threatening Behaviour, 33(2): 201-10.
54051	Chan C, Shaw T (2009). Suicidality induced by steroid treatment. Aust N
34031	Z J Psychiatry, 43(5): 484-5.
53985	Cheah D, Schmitt G, Pridmore S (2008). Suicide, misappropriation and
00000	impulsivity. Aust N Z J Psychiatry, 42: 544-6.
	Chen J, Dunne MP, Han P (2006). Child sexual abuse in Henan
52466	province, China: associations with sadness, suicidality, and risk
	behaviors among adolescent girls. J Adolesc Health, 38: 544-9.
69521	Chen LP, Murad MH, Paras ML, et al (2010). Sexual abuse and lifetime
	diagnosis of psychiatric disorders: systematic review and meta-analysis.
	Mayo Clin Proc, 85(7): 618-29.  Chang AT (1995) Montal illness and suicide: a case central study from
1744	Cheng AT (1995). Mental illness and suicide: a case-control study from
	east Taiwan. Arch Gen Psychiatry, 52(7): 594-603.  Cheng ATA, Chen THH, Chen CC, (2000). Psychosocial and psychiatric
54739	risk factors for suicide. Br J Psychiatry, 177: 360-5.
	There is the suicide. Did i sychiatry, 177. 000-0.

54325	Cheung A (2009). [Comment] Review: SSRIs are associated with increased risk for attempted or completed suicide in adolescents but not in adults or the elderly. ACP Journal Club, 150(6): JC6-13. Comment on ID: 54323.
52459	Cheung YB, Law CK, Chan B, et al (2006). Suicidal ideation and suicidal attempts in a population-based study of Chinese people: Risk attributable to hopelessness, depression, and social factors. Journal of Affective Disorders, 90: 193-9.
54730	Chia B-H, Chia A, Tai B-C (2008). Suicide letters in Singapore. Arch Suicide Res, 12: 74-81.
77602	Chou W-J, Liu T-L, Hu H-F, et al (2016). Suicidality and its relationships with individual, family, peer, and psychopathology factors among adolescents with attention-deficit/hyperactivity disorder. Res Dev Disabil, 53-54: 86-94.
48438	Choudary PV, Knowles JA (2008). Suicide and Impulsive Behaviour. Textbook of Psychiatry, 5th Edition, Part II, Chapter 6. American Psychiatric Publishing, Inc.
54205	Chowdhury AN, Banerjee S, Brahma A, et al (2007). Pesticide practices and suicide among farmers of the Sundarban region in India. Food & Nutrition Bulletin, 28(Suppl 2): S381-91.
77801	Christiansen E, Agerbo E, Bilenberg N, et al (2016). SSRIs and risk of suicide attempts in young people- a Danish observational register-based historical cohort study, using propensity score. Nord J Psychiatry, 70(3): 167-75.
52966	Cipriani A, Pretty H, Hawton K, et al (2005). Lithium in the prevention of suicidal behavior and all-cause mortality in patients with mood disorders: a systematic review of randomised trials. Am J Psychiatry, 162: 1805-19.
52450	Clarke DE, Colantonio A, Rhodes AE, et al (2008). Pathways to suicidality across ethnic groups in Canadian adults: the possible role of social stress. Psychological Medicine, 38: 419-31.
52160	Clarke VA, Frankish CJ, Green LW (1997). Understanding suicide among indigenous adolescents: a review using the PRECEDE model. Inj Prev, 3: 126-34.
52665	Clayton PJ (1985). Suicide. Psychiatr Clin N Am, 8(2): 203-14.
9664	Cohen BM, Cooper MZ (1954). A Follow-up Study of World War II Prisoners Of War, . US VA Medical Monograph, US Government Printing Office.
54682	Cole DA, Martin JM, Powers B, et al (1996). Modelling causal relations between academic and social competence and depression: a multitrait-multimethod longitudinal study of children. Journal of Abnormal Psychology, 105(2): 258-70.
52156	Colucci E, Martin G (2007). Ethnocultural aspects of suicide in young people: a systematic literature review part 2: risk factors, precipitating agents, and attitudes toward suicide. Suicide & Life-Threatening Behaviour, 37(2): 222-37.
85892	Compton S, Levy P, Griffin M, et al (2011). Family-witnessed resuscitation: Bereavement outcomes in an urban environment. Journal of Palliative Medicine, 14(6): 715-21.
54693	Conaghan S, Davidson KM (2002). Hopelessness and the anticipation of positive and negative future experiences in older parasuicidal adults. Br J Clin Psychol, 41: 233-42.
52464	Consolacion TB, Russell ST, Sue S (2004). Sex, race/ethnicity, and romantic attractions: multiple minority status adolescents and mental health. Cultur Divers Ethnic Minor Psychol, 10(3): 200-14.

79754	Coope C, Donovan J, Wilson C, et al (2015). Characteristics of people
	dying by suicide after job loss, financial difficulties and other economic
	stressors during a period of recession (2010-2011): a review of coroners'
54741	records. J Affect Disord, 183: 98-105.
	Cooper J, Kapur N, Webb R, et al (2005). Suicide after deliberate self-
	harm: a 4-year cohort study. Am J Psychiatry, 162: 297-303.
4400	Coryell W, Noyes R, Clancy J (1982). Excess mortality in panic disorder:
4188	A comparison with primary unipolar depression. Arch Gen Psychiatry, 39:
	701-3.  Coryell W, Noyes R, House JD (1986). Mortality among outpatients with
4186	anxiety disorders. Am J Psychiatry, 143(4): 508-10.
	Coupland C, Hill T, Morriss R, et al (2015). Antidepressant use and risk
77613	of suicide and attempted suicide or self harm in people aged 20 to 64:
77013	cohort study using a primary care database. BMJ, 350: h517.
	Coupland CAC, Dhiman P, Barton G, et al (2011). A study of the safety
77615	and harms of antidepressant drugs for older people: a cohort study using
77010	a large primary care database. Health Technology Assessment, 15(28).
	Cox BJ, Direnfeld DM, Swinson RP, et al (1994). Suicidal ideation and
4187	suicide attempts in panic disorder and social phobia. Am J Psychiatry,
1.0.	151(6): 882-7.
54045	Crombie IK (2001). [Comment] An interesting finding, but what does it
54015	mean? BMJ, 323: 5.
	Crow SJ, Peterson CB, Swanson SA,et al (2009). Increased mortality in
55079	bulimia nervosa and other eating disorders. Am J Psychiatry, : [Epub
	ahead of print].
4160	Crumley F E (1989). PTSD and suicide attempts by adolescents. Am J
4169	Psychiatry, 146(4): 558.
	Cukor J, Wyka K, Jayasinghe N, et al (2011). Prevalence and predictors
84177	of posttraumatic stress symptoms in utility workers deployed to the World
04177	Trade Center following the attacks of September 11, 2001. Depress
	Anxiety, 28(3): 210-7.
	Cunnigham FE, Hur K, Dong D, et al (2016). A comparison of
77479	neuropsychiatric adverse events during early treatment with varenicline
	or a nicotine patch. Addiction, (Epub ahead of print).
77475	Czabanski A, Lester D (2013). Suicide among Polish officers during
	World War II in Oflag II-C Woldenberg. Psychol Rep, 112(3): 727-31.
E 4075	Dahlen ER, Canetto SS (2002). The role of gender and suicide
54675	precipitant in attitudes toward nonfatal suicidal behavior. Death Studies,
	26: 99-116.
05011	Dai W, Chen L, Lai Z, et al (2016). The incidence of post-traumatic stress
85911	disorder among survivors after earthquakes: a systematic review and
	meta-analysis. BMC Psychiatry, 16: 188.  Daniel AE, Fleming J (2005). Serious suicide attempts in a state
54681	correctional system and strategies to prevent suicide. The Journal of
34001	Psychiatry and Law, 33: 227-47.
	Darke S, Kaye S, McKetin R, (2008). Major physical and psychological
54267	harms of methamphetamine use. Drug and Alcohol Review, 27: 253-62.
	Davidson JRT, Hughes D, Blazer DG, et al (1991). Post-traumatic stress
4171	disorder in the community: An epidemiological study. Psychological
7171	Medicine, 21: 713-21.
1745	Davis AT, Kosky RJ (1991). Attempted suicide in Adelaide and Perth:
	changing rates for males and females, 1971-1987. MJA, 154: 666-70.
	Davison EH, Kaiser AP, Spiro A, et al (2016). From late-onset stress
79752	symptomatology to later-adulthood trauma reengagement in aging
- 3-	combat veterans: taking a broader view. The Gerontologist, 56(1): 14-21.
	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1

54040	De Filippis S, Erbuto D, Gentili F, et al (2008). Mental turmoil, suicide
54010	risk, illness perception, and temperament, and their impact on quality of
	life in chronic daily headache. J Headache Pain, 9: 349-57.
54684	De Leo D (1996). Dementia, insight, and suicidal behavior. Crisis, 17(4): 147-8.
54687	De Leo D, Hickey PA, Meneghel G, et al (1999). Blindness, fear of sight
0-1001	loss, and suicide. Psychosomatics, 40(4): 339-44.
54831	De Leo D, Pellegrini C, Serraiotto L (1986). Adjustment disorders and suicidality. Psychol Rep, 59: 355-8.
52150	De Leo D, Spathonis K (2003). Suicide and euthanasia in late life. Aging Clin Exp Res, 15: 99-110.
	de Oliveira Ribeiro NP, de Mello Schier AR, Ornelas AC, et al (2013).
77472	Anxiety, depression and suicidal ideation in patients with rheumatoid
11412	arthritis in use of methotrexate, hydroxychloroquine, leflunomide and
	biological drugs. Compr Psychiatry, 54(8): 1185-9.
	Degenhardt L, Hall W, Warner-Smith M (2006). Using cohort studies to
54049	estimate mortality among injecting drug users that is not attributable to
	AIDS. Sex Transm Infect, 82(Suppl III): iii56-63.
5.47.40	Department of Health and Ageing (2007). Life. Research and evidence in
54742	suicide prevention. Retrieved 12 November 2009, from
	http://www.livingisforeveryone.com.au
77693	Devries KM, Mak JYT, Child JC, et al (2014). Childhood sexual abuse
	and suicidal behavior a meta-analysis. Pediatrics, 133: e1331-44.
54720	Dharmani C, Jaga K (2005). Epidemiology of acute organophosphate
54720	poisoning in hospital emergency room patients. Rev Environ Health, 20(3): 215-32.
	Diekstra RF, Garnefski N (1995). On the nature, magnitude, and
1747	causality of suicidal behaviors: an international perspective. Suicide Life
1777	Threat Behav, 25(1): 36-57.
	Diekstra RF, Gulbinat W (1993). The epidemiology of suicidal behaviour:
1748	a review of three continents. Wld Hlth Statist Quart, 46: 52-68.
	Dieperink E, Ho SB, Thuras P, et al (2003). A prospective study of
54322	neuropsychiatric symptoms associated with intefern-a-2b and ribavirin
	therapy for patients with chronic hepatitis C. Psychosomatics, 44: 104-12.
	Dieperink E, Ho SB, Tetrick L, et al (2004). Suicidal ideation during
53450	interferon-a2b and ribavirin treatment of patients with chronic hepatitis C.
	General Hospital Psychiatry, 26: 237-40.
	Dilsaver SC, Benazzi F, Akiskal KK, et al (2008). Differential patterns of
53475	lifetime multiple anxiety disorder comorbidity between Latino adults with
00470	bipolar I and major depressive disorders. Bulletin of the Menninger Clinic,
	72(2): 130-48.
4174	Dilsaver SC, Chen YW, Swann AC, et al (1994). Suicidality in patients
	with pure and depressive mania. Am J Psychiatry, 151(9): 1312-5.
77929	Dixon JB (2016). Self-harm and suicide after bariatric surgery: time for
	action. Lancet Diabetes Endocrinol, 4(3): 199-200.
05040	Dobashi K, Nagamine M, Shigemura J, et al (2014). Psychological effects
85912	of disaster relief activities on Japan ground self-defense force personnel following the 2011 great east Japan earthquake. Psychiatry, 77: 190-9.
	Dobry Y, Dolores M, Sher B, et al (2013). Bullying, psychiatric pathology
77367	and suicidal behavior. Int J Adolesc Med Health, 25(3): 295-9.
	Duberstein PR, Conwell Y, Conner KR, et al (2004). Suicide at 50 years
54767	of age and older: perceived physical illness, family discord and financial
	strain. Psychol Med, 34: 137-46.
	Dunt D (2009). Independent study into suicide in the ex-service
54294	community Retrieved 25 September 2009, from www.dva.gov.au
50450	Eastwood MR, Peacocke JE (1975). Suicide, diagnosis and age. Can
52152	Psychiatr Assoc J, 20: 447-9.

F.4000	Eddleston M (2008). [Comment] The pathophysiology of
54208	organophosphorus pesticide self-poisoning is not so simple. The
	Netherlands Journal of Medicine, 66(4): 146-8. Comment on ID: 54207.
F400F	Edwards RR, Smith MT, Kudel I, et al (2006). Pain-related
54005	catastrophizing as a risk factor for suicidal ideation in chronic pain. Pain,
	126: 272-9. Eisenberg ME, Neumark-Sztainer D, Story M (2003). Associations of
52726	weight-based teasing and emotional well-being among adolescents. Arch
53726	Pediatr Adolesc Med, 157: 733-8.
	Elklit A, Kurdahl S (2013). The psychological reactions after witnessing a
85881	killing in public in a Danish high school. European Journal of
03001	Psychotraumatology, 4: 1-7.
	Elmes NJ, Nasveld PE, Kitchener SJ, et al (2008). The efficacy and
	tolerability of three different regimens of tafenoquine versus primaquine
78018	for post-exposure prophylaxis of plasmodium vivax malaria in the
	southwest pacific. Trans R Soc Trop Med Hyg, 102(11): 1095-101.
70047	Engel CC (2013). Suicide, mental disorders, and the US military: Time to
79647	focus on mental health service delivery. JAMA, 310(5): 484-5.
	Epstein RS, Fullerton CS, Ursano RJ (1998). Posttraumatic stress
86474	disorder following an air disaster: a prospective study. Am J Psychiatry,
	155(7): 934-8.
	Erlangsen A, Stenager E, Conwell Y (2015). Physical diseases as
77763	predictors of suicide in older adults: a nationwide, register-based cohort
	study. Soc Psychiatry Psychiatr Epidemiol, 50: 1427-39.
	Espelage DL, Holt MK (2013). Suicidal ideation and school bullying
77372	experiences after controlling for depression and delinquency. J Adolesc
	Health, 53: S27-S31.
	Espie E, Gaboulaud V, Baubet V, et al (2009). Trauma-related
85904	psychological disorders among Palestinian children and adults in Gaza
	and West Bank, 2005-2008. International Journal of Mental Health
	Systems, 3(21): 1-5.
52674	Evans DL (1982). Explaining suicide among the young: An analytical review of the literature. JPNMHS, 20(8): 9-16.
	Evans E, Hawton K, Rodham K, Deeks J (2005). The prevalence of
52462	suicidal phenomena in adolescents: A systemic review of population-
02102	based studies. Suicide & Life-Threatening Behaviour, 35(3): 239-50.
	Evans E, Hawton K, Rodham K, et al (2005). The prevalence of suicidal
52677	phenomena in adolescents: A systemic review of population-based
0_0.1	studies. Suicide & Life-Threatening Behaviour, 35(3): 239-50.
	Fang F, Keating NL, Mucci LA, et al (2010). Immediate risk of suicide and
77609	cardiovascular death after a prostate cancer diagnosis: cohort study in
	the United States. J Natl Can Inst, 102: 307-14.
	Fanous AH, Prescott CA, Kendler KS (2004). The prediction of thoughts
52441	of death or self-harm in a population-based sample of female twins.
	Psychological Medicine, 34: 301-12.
	Farberow NL, Kang HK, Bullman TA (1990). Combat experience and
4168	postservice psychosocial status as predictors of suicide in Vietnam
	veterans. The Journal of Nervous and Mental Disease, 38(1): 32-7.
76956	Fardet L, Petersen I, Nazareth I (2012). Suicidal behavior and severe
	neuropsychiatric disorders following glucocorticoid therapy in primary
	care. Am J Psychiatry, 169(5): 491-7.
52161	Farrer LA (1986). Suicide and attempted suicide in Huntington disease: implications for preclinical testing of persons at risk. Am J Med Genet, 24:
	305-11.
	Fassberg MM, van Orden KA, Duberstein P, et al (2012). A systematic
77373	review of social factors and suicidal behavior in older adulthood. Int J
1,070	Environ Res Public Health, 9: 722-45.
	Entra entre en abaie riodail, o. 122 io.

52443	Fathelrahman AI, Ab Rahman AF, Mohd Zain Z (2008). Self-poisoning by drugs and chemicals: variations in demographics, associated factors and
52151	final outcomes. General Hospital Psychiatry, 30: 467-70.  Fazel S, Cartwright J, Norman-Nott A, et al (2008). Suicide in prisoners: a systematic review of risk factors. J Clin Psychiatry, 69: 1721-31.
77618	Fazel S, Wolf A, Pillas D, et al (2014). Suicide, fatal injuries, and other causes of premature mortality in patients with traumatic brain injury: a 41-year Swedish population study. JAMA Psychiatry, 71(3): 326-33.
54217	FDA (2004). Drugs. Suicidality in children and adolescents being treated with antidepressant medications. Retrieved 15 September 2009, from http://www.fda.gov/Drugs/DrugSafety/PublicHealthAdvisories/ucm161679.htm
54218	FDA (2009). Drugs. Strattera. Public health advisory - suicidal thinking in children and adolescents being treated with Strattera (atomoxetine). Retrieved 21 September 2009, from http://www.fda.gov/Drugs/DrugSafety/PublicHealthAdvisories/ucm051733.htm
77979	FDA (2013). FDA Drug Safety Communication: FDA approves label changes for antimalarial drug mefloquine hydrochloride due to risk of serious psychiatric and nerve side effects. Retrieved 15 April 2016, from http://www.fda.gov/drugs/drugsafety/icm362227.htm
77928	FDA (2013). Information for healthcare professionals: atomoxetine (marketed as strattera). Retrieved 6 April 2016, from http://www.fda.gov/Drugs/DrugSafety/PostmarketDrugSafetyInformationf orPatientsandProviders/ucm124391.htm
77631	FDA (2014). Memorandum - overview of the October 16, 2014 joint meeting of the psychopharmacologic drugs advisory committee and drugs safety and risk management advisory committee meeting. FDA Briefing Document, 4-25.
77554	FDA Advisory Committee (2014). Executive Summary Only. Joint meeting of the psychopharmacologic drugs advisory committee and the drug safety and risk management advisory committee: risk of serious neuropsychiatric adverse events with chantix (varenicline tartrate). Pfizer.
54216	FDA U.S. Food and Drug Administration (2007). FDA proposes new warnings about suicidal thinking, behavior in young adults who take antidepressant medication. Retrieved 15 September 2009, from http://www.fda.gov/NewsEvents/Newsroom/PressAnnouncements/2007/ucm108905.htm
54047	Fergusson DM, Horwood LJ, Beautrais AL (1999). Is sexual orientation related to mental health problems and suicidality in young people? Arch Gen Psychiatry, 56: 876-80.
85913	Ferry F, Bunting B, Murphy S, et al (2014). Traumatic events and their relative PTSD burden in Northern Ireland: a consideration of the impact of the 'Troubles'. Social Psychiatry & Psychiatric Epidemiology, 49(3): 435-46.
77843	Finley EP, Bollinger M, Noel PH, et al (2015). A national cohort study of the association between the polytrauma clinical triad and suicide-related behavior among US veterans who served in Iraq and Afghanistan. Am J Public Health, 105(2): 380-7.
54203	Fishbain DA (1999). The association of chronic pain and suicide. Semin Clin Neuropsych, 4(3): 221-7.
77555	Fishbain DA, Bruns D, Meyer LJ, et al (2012). Exploration of the relationship between disability perception, preference for death over disability, and suicidality in patients with acute and chronic pain. Pain Medicine, 13: 552-61.
54014	Fishbain DA, Bruns D, Disorbio JM, et al (2009). Risk for five forms of suicidality in acute pain patients and chronic pain patients vs pain-free community controls. Pain Medicine, [Epub ahead of print].

54013	Fishbain DA, Lewis JE, Gao J, et al (2009). Are chronic low back pain patients who smoke at greater risk for suicide ideation? Pain Medicine,
	10(2): 340-6.
77556	Fishbain DA, Lewis JE, Gao J (2014). The pain suicidality association: a narrative review. Pain Medicine, 15: 1835-49.
11032	Fitzgerald LF (1993). Sexual harassment. Violence against women in the
	workplace. American Psychologist, Vol 48(10): 1070-6.
	Fleischmann A, Bertolote JM, Belfer M, et al (2005). Completed suicide
54285	and psychiatric diagnoses in young people: a critical examination of the
0.1200	evidence. Am J Orthop, 75(4): 676-83.
	Flensborg-Madsen T, Knop J, Mortensen EL, et al (2009). Alcohol use
77610	disorders increase the risk of completed suicide- irrespective of other
	psychiatric disorders. A longitudinal cohort study. Psychiatry Res, 167:
	123-30.
	Fontana A, Rosenheck R (1995). Attempted suicide among Vietnam
1751	veterans: a model of etiology in a community sample. Am J Psychiatry,
	152(1): 102-9.
	Fontana A, Rosenheck R (1995). An etiological model of attempted
4173	suicide among Vietnam theater veterans: Prospective generalization to a
	treatment-seeking sample. J Nerv Ment Dis, 183: 377-83.
	Foreign Affairs, Defence and Trade References Committee (2017). The
83048	Constant Battle: Suicide by Veterans. Commonwealth of Australia.
50070	Fortuna LR, Perez DJ, Canino G, et al (2007). Prevalence and correlates
52670	of lifetime suicidal ideation and suicide attempts among Latino subgroups
	in the United States. J Clin Psychiatry, 68(4): 572-81.
	Fragoso YD, Frota ER, Lopes JS et al (2010). Severe depression, suicide
72582	attempts and ideation during the use of interferon beta by patients with
	multiple sclerosis. Clinical Neuropharmacology, 33(6): 312-6.
77842	Fralick M, Thiruchelvam D, Tien HC, et al (2016). Risk of suicide after a
11042	concussion. CMAJ, [Epub ahead of print].
	Frans O, Rimmo P-A, Aberg L, et al (2005). Trauma exposure and post-
34537	traumatic stress disorder in the general population. Acta Psychiatr Scand,
	111: 291-9.
	Fraser CE, Smith KB, Judd F, et al (2005). Farming and mental health
53997	problems and mental illness. Int J Soc Psychiatry, 51: 340-9.
	Freed G, Stringer PB (1968). Comparative mortality experience 1946-
13054	1963 among former Australian prisoners of war of the Japanese. Medical
13034	
	Research Bulletin, No 2. Central Medical Research Advisory Committee.
49920	Friedman H (1997). [Comment] Spinal cord injury questions. Mil Med,
	162(5): iii-iv.
	Friedman M, Glasser M, Laufer E, et al (1972). Attempted suicide and
54830	self-mutilation in adolescence: some observations from a psychoanalytic
	research group. Int J Psychoanal, 53: 179-83.
83834	Friedman MJ (2015). Risk factors for suicide among army personnel.
03034	JAMA, 313(11): 1154-5.
F000F	Friedman RC (1999). [Comment] Homosexuality, psychopathology, and
53685	suicidality. Arch Gen Psychiatry, 56: 887-8. Comment on ID: 53682.
4180	Friedman RC, Aronoff MS, Clarkin JF, et al (1983). History of suicidal
	behaviour in depressed borderline inpatients. Am J Psychiatry, 140(8):
	1023-6.
	Friedman S, Jones JC, Cherner L, et al (1992). Suicidal ideation and
4200	suicide attempts among patients with panic disorder: A survey of two
	outpatients clinics. Am J Psychiatry, 149(5): 680-5.
53503	Frisman LK, Griffin-Fennell F (2009). [Comment] Suicide and
	incarcerated veterans - don't wait for the numbers. J Am Acad Psychiatry
	Law, 37: 92-4. Comment on ID: 53500.

	- ND
4170	Fryer MR, Frances AJ, Sullivan T, et al (1988). Suicide attempts in
4179	patients with borderline personality disorder. Am J Psychiatry, 145(6):
	737-9.
53995	Fuller JD, Kelly B, Law S, et al (2009). Service network analysis for
	agricultural mental health. BMC Health Services Research, 9: 87.
69181	Fullerton CS, Ursano RJ, Reeves J, et al (2006). Perceived safety in
-	disaster workers following 9/11. J Nerv Ment Dis, 194(1): 61-3.
52664	Galanter M, Castaneda R (1985). Self-destructive behavior in the
	substance abuser. Psychiatr Clin N Am, 8(2): 251-61.
04070	Gale CR, Braidwood EA, Winter PD, et al (1999). Mortality from
21879	Parkinson's disease and other causes in men who were prisoners of war
	in the Far East. The Lancet, 354(9196): 2116-8.
E2002	Gallagher LM, Kliem C, Beautrais AL, et al (2008). Suicide and
53993	occupation in New Zealand, 2001-2005. Int J Occup Environ Health, 14:
	45-50.
1752	Gallerani M, Manfredini R, Caracciolo S, et al (1995). Serum cholesterol
	concentrations in parasuicide. BMJ, 310(6995): 1632-6.  Gardner DL, Cowdry RW (1985). Suicidal and parasuicidal behaviour in
52662	
	borderline personality disorder. Psychiatr Clin N Am, 8(2): 389-403.  Garlow SJ, Purselle D, D'Orio B (2008). Nonpharmacological Psychiatric
48437	Emergencies. Essentials of Clinical Psychopharmacology, Second
40437	Edition, Chapter 41. American Psychiatric Publishing, Inc.
_	Geddes JR, Barbui C, Cipriani A (2009). Risk of suicidal behaviour in
	adults taking antidepressants. Increased risk is probably restricted to
54024	younger people and varies greatly between individual medicines. BMJ,
	339: 411-2.
	Gerson J, Stanley B (2002). Suicidal and self-injurious behavior in
45829	personality disorder: controversies and treatment directions. Current
40020	Psychiatry Reports, 4: 30-8.
_	Geulayov G, Gunnell D, Holmen TL, et al (2012). The association of
	parental fatal and non-fatal suicidal behaviour with offspring suicidal
78035	behaviour and depression: a systematic review and meta-analysis.
	Psychol Med, 42(8): 1567-80.
	Ghaemi SN, Vohringer PA, Whitham EA (2013). Antidepressants from a
77486	public health perspective: re-examining effectiveness, suicide, and
	carcinogenicity. Acta Psychiatr Scan, 127(2): 89-93.
_	Gibbons R, Mann JJ (2009). [Comment] Proper studies of selective
54324	serotonin reuptake inhibitors are needed for youth with depression.
	CMAJ, 180(3): 270-1. Comment on ID: 54323.
77557	Gibbons RD, Mann JJ (2013). Varenicline, smoking cessation and
77557	neuropsychiatric adverse events. Am J Psychiatry, 170(12): 1460-7.
77507	Glasheen C, Pemberton MR, Lipari R, et al (2015). Binge drinking and
77597	the risk of suicidal thoughts, plans, and attempts. Addict Behav, 43: 42-9.
	Gnanarai P, Karthikevan S, Narasimhan M, et al (2015). Decrease in
78062	"Hamilton rating scale for depression" following isotretinoin therapy in
	acne: an open-label prospective study. Indian J Dermatol, 60(5): 461-4.
54031	Goeb JL, Even C, Nicolas G, et al (2006). Psychiatric side effects of
J <del>4</del> UJ I	interferon-B in multiple sclerosis. Eur Psychiatry, 21: 186-93.
11033	Golding JM (1994). Sexual assault history and physical health in
11000	randomly selected Los Angeles women. Health Psychology, 13(4): 130-8.
	Goldston DB, Daniel SS, Erkanli A, et al (2009). Psychiatric diagnoses as
77766	contemporaneous risk factors for suicide attempts among adolescents
77700	and young adults: developmental changes. J Consult Clin Psychol, 77(2):
	281.
1753	Golier JA, Marzuk PM, Leon AC, et al (1995). Low serum cholesterol
	level and attempted suicide. Am J Psychiatry, 152: 419-23.

77558	Gordon M, Melvin G (2013). Selective serotonin re-uptake inhibitors: a review of the side effects in adolescents. Aust Fam Physician, 42(9): 620-
77559	3. Gorton HC, Webb RT, Kapur N, et al (2016). Non-psychotropic medication and risk of suicide or attempted suicide a systematic review. BMJ, 6: e009074.
77605	Gradus JL, Antonsen S, Svensson E, et al (2015). Trauma, comorbidity, and mortality following diagnoses of severe stress and adjustment disorders: a nationwide cohort study. Am J Epidemiol, 182(5): 451-8.
77619	Gradus JL, Qin P, Lincoln AK, et al (2010). Acute stress reaction and completed suicide. Int J Epidemiol, 39: 1478-84.
9565	Green BL (1990). Defining trauma: terminology and generic stressor dimensions. J Applied Social Psychology, 20: 1632-42.
78013	Green JA, Patel AK, Patel BR, et al (2014). Tafenoquine at therapeutic concentrations does not prolong fridericia-corrected QT interval in healthy subjects. J Clin Pharmacol, 54(9): 995-1005.
85636	Gross R, Neria Y, Tao XG, et al (2006). Posttraumatic stress disorder and other psychological sequelae among World Trade Center clean up and recovery workers. Ann N Y Acad Sci, 1071: 495-9.
52159	Grossman D (1997). [Comment] Watching the canary: the prevention of suicide. Inj Prev, 3: 74-7.
52149	Groves SA, Stanley BH, Sher L (2007). Ethnicity and the relationship between adolescent alcohol use and suicidal behaviour. Int J Adolesc Med Health, 19(1): 19-25.
77129	Grupp D, Froscher RW (1994). (article in German) Neuropsychiatrishe storungen nach malariaprophylaxe mit mefloquine. Eng - Neuropsychiatric disturbances after malaria prophylaxis with mefloquine. Akt Neurol, 21: 134-6.
8785	Guest CS, Venn AJ (1992). Mortality of former prisoners of war and other Australian veterans. Med J Aust, 157(2): 132-5.
77982	Gulgin M-B, Li J, Pedersen HS, et al (2015). Incidence of suicide among persons who had a parent who died during their childhood: a population-based cohort study. JAMA Psychiatry, 72(12): 1227-34.
53687	Gutierrez PM, Brenner LA, Huggins JA (2008). A preliminary investigation of suicidality in psychiatrically hospitalized veterans with traumatic brain injury. Arch Suicide Res, 12: 336-43.
78021	Hale BR, Owusu-Agyei S, Fryauff DJ, et al (2003). A randomized, double-blind, placebo-controlled, dose-ranging trial of tafenoquine for weekly prophylaxis against plasmodium falciparum. Clin Infect Dis, 36(5): 541-9.
9449	Hall RCW, Malone PT (1976). Psychiatric effects of prolonged Asian captivity: A two-year follow-up. Am J Psychiatry, 133(7): 786-90.
39810	Hammad TA, Laughren TP, Racoosin JA (2006). Suicide rates in short-term randomized controlled trials of newer antidepressants. Journal of Clinical Psychopharmacology, 26(2): 203-7.
11034	Hampton H (1995). Care of the woman who has been raped. NEJM, 332(4): 234-7.
79679	Handley TE, Inder KJ, Key-Lambkin FJ, et al (2012). Contributors to suicidality in rural communities beyond the effects of depression. BMC Psychiatry, 12: 105.
54572	Harrison JE, Pointer S, Elnour AA (2009). A review of suicide statistics in Australia. Injury Research and Statistics Series, Number 49. Australian Institute of Health and Welfare, Canberra.
84414	Hatton AT (2011). The experience of witnessing a stranger's suicide [Excerpt from book]. Dissertation Abstracts International: Section B: The Sciences and Engineering, 74(1-B {E}) [Abstract]

49088	Haukka J, Tiihonen J, Harkanen T, et al (2008). Association between medication and risk of suicide, attempted suicide and death in nationwide cohort of suicidal patients with schizophrenia. Pharmacoepidemiology and Drug Safety, 17: 686-96.
54500	Hawgood J, De Leo D (2008). Anxiety disorders and suicidal behaviour: an update. Curr Opin Psychiatry, 21: 51-64.
73351	Hawthorne G, Korn S, Creamer M (2014). Australian peacekeepers: Long-term mental health status, health service use, and quality of life. Summary Report. Australian Centre for Posttraumatic Mental Health, Department of Psychiatry, The University of Melbourne.
1754	Hawton K, Fagg J (1988). Suicide, and other causes of death, following attempted suicide. Br J Psychiatry, 152: 359-66.
1755	Hawton K, Fagg J, Platt S, et al (1993). Factors associated with suicide after parasuicide in young people. BMJ, 306: 1641-4.
54738	Hawton K, Harriss L (2006). Deliberate self-harm in people ages 60 years and over: characteristics and outcome of a 20-year cohort. Int J Geriatr Psychiatry, 21: 571-81. Retrieved 12 November 2009, from 0
54685	Hayes LM (2001). Jail suicide risk despite denial (or: when actions speak louder than words). Crisis, 22(1): 7-9.
54018	Healy D, Whitaker C (2003). Antidepressants and suicide: risk-benefit conundrums. J Psychiatry Neurosci, 28(5): 331-7.
77560	Hejri SM, Faizi M, Babaeian M (2013). Zolpidem-induced suicide attempt: a case report. DARU Journal of Pharmaceutical Sciences, 21: 77.
54727	Heller TS, Hawgood JL, De Leo D (2007). Correlates of suicide in building industry workers. Arch Suicide Res, 11: 105-17.
4170	Hendin H, Haas A P (1991). Suicide and guilt as manifestations of PTSD in Vietnam combat veterans. Am J Psychiatry, 148(5): 586-91.
83743	Hepinstall E, et al (2004). PTSD and depression in refugee children: associations with pre-migration trauma and post-migration stress. European Child and Adolescent Psychiatry, 13(6): 373-80.
53682	Herrell R, Goldberg J, True WR, et al (1999). Sexual orientation and suicidality. A co-twin control study in adult men. Arch Gen Psychiatry, 56: 867-74.
77723	Hesdorffer DC, Ishihara L, Webb DJ (2016). Occurrence and recurrence of attempted suicide among people with epilepsy. JAMA Psychiatry, 73(1): 80-6.
54026	Hesdorffer DC, Kanner AM (2009). The FDA alert of suicidality and antiepileptic drugs: fire or false alarm? Epilepsia, 50(5): 978-86.
77616	Hetrick S (2016). Monitoring of suicide risk throughout the course of treatment with antidepressants for depression is required, but vigilance is required for those on some particular antidepressant agents. Evid Based Ment Health, 18(3): 86.
77815	Hetrick SE, McKenzie JE, Cox GR, et al (2012). Newer generation antidepressants for depressive disorders in children and adolescents (review). Cochrane Database of Systematic Reviews, (11): CD004851.
54809	Hierholzer R (2001). Suicide in dementia: case studies of failure as a risk factor. Clinical Gerontologist, 24(1-2): 159-63.
54692	Hill RD, Gallagher D, Thompson LW, et al (1988). Hopelessness as a measure of suicidal intent in the depressed elderly. Psychology and Aging, 3(3): 230-2.
54688	Hine TJ, Pitchford NJ, et al (2000). [Comments] Blindness and high suicide risk? Psychosomatics, 41(4): 370-1. Comments on ID: 54687.
77848	Holla S, Amberkar MB, Bhandarypanambur R, at el (2015). Cycloserine induced late onset psychosis and ethambutol induced peripheral neuropathy associated with MDR-TB treatment in an Indian patient- a rare case report. Journal of Clinical and Diagnostic Research, 9(2): FD01-3.

r	
77374	Holt MK, Vivolo-Kantor AM, Polanin JR, et al (2015). Bullying and suicidal ideation and behaviors: a meta-analysis. Pediatrics, 135(2): e496-e509.
77476	Hooley JM, Franklin JC, Nock MK (2014). Chronic pain and suicide: understanding the association. Curr Pain Headache Rep, 18(8): 435.
4182	Hornig CD, McNally RJ (1995). Panic disorder and suicide attempt: A reanalysis of data from the epidemiologic catchment area study. British Journal of Psychiatry, 167: 76-9.
85400	Huang J, Liu Q, Li X, et al (2013). Post-traumatic stress disorder status in a rescue group after the Wenchuan earthquake relief. Neural Regen Res, 8(20): 1898-1906.
4193	Hughes M (1994). Cholesterol and violent death: Diets, violence and civilisation. BMJ, 309: 1228.
52158	Hunter E, Harvey D (2002). Indigenous suicide in Australia, New Zealand, Canada and the United States. Emerg Med, 14: 14-23.
52162	Huong TTT, Szentesi I, Czeizel A (1988). Lower prevalence of chromosome aberrations and SCEs in self-poisoned pregnant women. Mutation Research, 198: 255-9.
71886	Ikin JF, McKenzie DP, Creamer MC et al (2005). War zone stress without direct combat: the Australian naval experience of the Gulf War. Journal of Traumatic Stress, 18(3): 193-204.
32984	Ikin JF, Sim MR, Creamer MC, et al (2004). War-related psychological stressors and risk of psychological disorders in Australian veterans of the 1991 Gulf War. British Journal of Psychiatry, 185: 116-26.
77826	Ilgen MA, Bohnert ASB, Ganoczy D, et al (2016). Opioid dose and risk of suicide. Pain, [Epub ahead of print]: DOI:10.1097/j.pain.000000000000484.
54012	Ilgen MA, Zivin K, McCammon RJ, et al (2008). Pain and suicidal thoughts, plans and attempts in the United States. General Hospital Psychiatry, 30: 521-7.
77604	Impey M, Heun R (2012). Completed suicide, ideation and attempt in attention deficit hyperactivity disorder. Acta Psychiatr Scand, 125(2): 93-102.
54683	Ingoldsby EM, Kohl GO, McMahon RJ, et al (2006). Conduct problems, depressive symptomatology and their co-occurring presentation in childhood as predictors of adjustment in early adolescence. J Abnorm Child Psychol, 34: 603-21.
54043	Inoue K (2009). [Comment] Significant correlation of the change in the divorce rate with the suicide rate in Japan from 1992 to 2004. Am J Forensic Med Pathol, 30(3): 311-2.
77849	Institute of Medicine (2008). Long-term consequences of traumatic brain injury. Gulf War and Health, Vol 7: 265-300. National Academies Press-Washington DC.
50832	Institute of Medicine of the National Academies (2008). Physiologic, Psychologic, and Psychosocial Effects of Deployment-Related Stress. Gulf War and Health, Volume 6. The National Academies Press, Washington, D.C.
73310	Irwig MS (2012). Depressive symptoms and suicidal thoughts among former users of finasteride with persistent sexual side effects. J Clin Psychiatry, 73(9): 1220-3.
11035	Isac M, Schneider S (1992). Some psychological reactions of rape victims. Medicine and Law, 11: 303-8.
77375	Isometsa E (2014). Suicidal behaviour in mood disorders - who, when, and why? Can J Psychiatry, 59(3): 120-30.
53673	Ivarsson T, Broberg AG, Arvidsson T, et al (2005). Bullying in adolescence: psychiatric problems in victims and bullies as measured by the Youth Self Report (YSR) and the Depression Self-Rating Scale (DSRA). Nordic Journal of Psychiatry, 59(5): 365-73.

54578	Jablonska B, Lindberg L, Lindblad F, et al (2009). School performance and hospital admissions due to self-inflicted injury: a Swedish national cohort study. Int J Epidemiol, 38: 1134-41.
45597	Jacobs DG, Deutsch NL, Brewer M (2001). Suicide, depression, and isotretinoin: is there a causal link? J Am Acad Dermatol, 45(5): S168-S175.
73105	Jacobson IG, Ryan MA, Hooper TI, et al (2008). Alcohol use and alcohol-related problems before and after military combat deployment. JAMA, 300: 663-75.
54270	Jaga K, Dharmani C (2007). The interrelation between organophosphate toxicity and the epidemiology of depression and suicide. Rev Environ Health, 22(1): 57-73.
54206	Jaga K, Dharmani C (2007). The interrelation between organophosphate toxicity and the epidemiology of depression and suicide. Reviews on Environmental Health, 22(1): 57-73.
78395	Jakobsen IS, Christiansen E (2011). Young people's risk of suicide attempts in relation to parental death: a population-based register study. Journal of Child Psychology and Psychiatry, 52(2): 176-83.
79751	Jakupcak M, Hoerster KD, Varra A, et al (2011). Hopelessness and suicidal ideation in Iraq and Afghanistan war veterans reporting subthreshold and threshold posttraumatic stress disorder. J Nerv Ment Dis, 199: 272-5.
71870	James LM, Van Kampen E, Miller RD et al (2013). Risk and protective factors associated with symptoms of post-traumatic stress, depression and alcohol misuse in OEF/OIF veterans. Military Medicine, 178(2): 159-65.
54689	Jensen JG, Jensen BF, Arensman E, et al, (1999). Attempted suicide and major public holidays in Europe: findings from the WHO/EURO multicentre study on parasuicide. Acta Psychiatr Scand, 99: 412-8.
54019	Jepsen P, Johnsen SP, Sorensen HT (2003). Risk of suicide in users of cardiovascular drugs. A review of the epidemiological evidence. Am J Cardiovasc Drugs, 3(3): 163-7.
54038	Jick H, Hagberg KW, Egger P (2009). Rate of suicide in patients taking montelukast. Pharmacotherapy, 29(2): 165-6.
39278	Jick SS, Kremers HM, Vasilakis-Scaramozza C (2000). Isotretinoin use and risk of depression, psychotic symptoms, suicide, and attempted suicide. Arch Dermatol, 136: 1231-36.
52469	Joe S, Baser RE, Breeden G, et al (2006). Prevalence of and risk factors for lifetime suicide attempts among blacks in the United States. JAMA, 296(17): 2112-23.
52155	Joe S, Niedermeier DM (2008). Social work research on African Americans and suicidal behaviour: a systematic 25-year review. Health & Social Work, 33(4): 249-57.
1756	Johnson J, Weissman MM, Klerman GL (1990). Panic disorder, comorbidity, and suicide attempts. Arch Gen Psychiatry, 47: 805-8.
53999	Johnson JG, Cohen P, Kasen S, et al (2002). Eating disorders during adolescence and the risk for physical and mental disorders during early adulthood. Arch Gen Psychiatry, 59: 545-52.
15070	Jordan BK, Schlenger WE, Hough R, et al (1991). Lifetime and current prevalence of specific psychiatric disorders among Vietnam Veterans and controls. Arch Gen Psychiatry, 48(3): 207-215.
75197	Jousset N, Rouge-Maillart C, Turcant A, et al (2010). Suicide by skull stab wounds. Am J Forensic Med Pathol, 31(4): 378-81.
53991	Judd F, Jackson H, Fraser C, et al (2006). Understanding suicide in Australian farmers. Soc Psychiatry Psychiatr Epidemiol, 41: 1-10.

	T 0.1 0.1 50 1/00/4/) 4.1
77482 54271	Judd LL, Schettler PJ, Brown ES, et al (2014). Adverse consequences of
	glucocorticoid medication: psychological, cognitive, and behavioral
	effects. Am J Psychiatry, 171(10): 1045-51.
	Juurlink DN, Herrmann N, Szalai JP, et al (2004). Medical illness and the risk of suicide in the elderly. Arch Intern Med, 164: 1179-84.
54370	Kaizar EE, Greenhouse JB, Seltman H, et al (2006). Do antidepressants
	cause suicidality in children? A Bayesian meta-analysis. Clinical Trials, 3:
	73-98.
	Kalant H (2001). The pharmacology and toxicology of "ecstasy" (MDMA)
54027	and related drugs. CMAJ, 165: 917-28.
	Kaltiala-Heino R, Rimpela M, Marttunen, et al (1999). Bullying,
53444	depression, and suicidal ideation in Finnish adolescents: school survey.
	BMJ, 319: 348-51.
	Kang HK, Bullman TA (2008). [Comment] Risk of suicide among US
53504	veterans after returning from the Iraq or Afghanistan war zones. JAMA,
	300(6): 652-3.
	Kang HK, Bullman TA, Smolenski DJ, et al (2015). Suicide risk among
81435	1,3 million veterans who were on active duty during the Iraq and
	Afghanistan wars. Ann Epidemiol, 25(2): 96-101.
	Kanner AM (2009). Depression and epilepsy: do glucocorticoids and
54268	glutamate explain their relationship? Curr Neurol Neurosci Rep, 9: 307-
	12.
50.400	Kaplan MS, Huguet N, McFarland BH, et al (2007). Suicide among male
53463	veterans: a prospective population-based study. J Epidemiol Community
	Health, 61: 619-24.
77944	Kaplan MS, Huguet N, McFarland BH, et al (2014). Use of alcohol before
	suicide in the United States. Ann Epidemiol, 24(8): 588-92.  Kaplan MS, McFarland BH, Huguet N, et al (2012). Suicide risk and
79680	precipitating circumstances among young, middle-aged, and older male
7 3000	veterans. Am J Public Health, 102: S131-7.
	Kaplan S, Lester D (1994). Depression, mania, and suicidal
4205	preoccupation. Psychological Reports, 74: 974.
50445	Kapur N, While D, Blatchley N, et al (2009). Suicide after leaving the UK
53445	armed forces - a cohort study. PLoS Med, 6(3): e1000026.
	Karch DL, Logan J, McDaniel DD, et al (2013). Precipitating
77376	circumstances of suicide among youth aged 10-17 years by sex: data
11310	from the national violent death reporting system, 16 states, 2005-2008. J
	Adolesc Health, 53: S51-S53.
78112	Karila L, Megarbane B, Cottencin O, et al (2015). Synthetic cathinones: a
70112	new public health problem. Current Neuropharmacology, 13: 12-20.
53449	Kausch O (2003). Suicide attempts among veterans seeking treatment
	for pathological gambling. J Clin Psychiatry, 64(9): 1031-8.
54679	Kausch O (2004). Irrational fear of AIDS associated with suicidal
	behavior. Journal of Psychiatric Practice, 10(4): 266-71.
	Kausch O and McCormick RA (2002). Suicide prevalence in chemical
28597	dependency programs: preliminary data from a national sample, and an
	examination of risk factors. Journal of Substance Abuse Treatment, 22: 97-102.
	Keane TM, King LA, King DW, et al (1998). Do war zone stressors
12622	predict the development of psychiatric disorders? RMA Conference
12022	Papers.
	Keane TM, Marshall AD, Taft CT (2006). Posttraumatic stress disorder:
69571	etiology, epidemiology, and treatment outcome. Annu Rev Clin Psychol,
	2: 161-97.
54828	Keck PE Jr, Merikangas KR, McElroy SL, et al (1994). Diagnostic and
	treatment implications of psychiatric comorbidity with migraine. Ann Clin
	Psychiatry, 6(3): 165-71.

	Keehn RJ (1980). Follow-up studies of World War II and Korean conflict
8923	prisoners: III. Mortality to January 1, 1976. Am J Epidemiol, 111: 194-211.
52663	Kellner CH, Best CL, Roberts JM, et al (1985). Self-destructive behavior in hospitalized medical and surgical patients. Psychiatr Clin N Am, 8(2): 279-89.
77698	Kennebeck S, Bonin L (2016). Suicidal behavior in children and adolescents: epidemiology and risk factors Retrieved 14 March 2016, from http://www.uptodate.com/contents/suicidal-behavior-in-children-and-adolescents-epidemiology-and-risk-factors
4213	Kerr TA, Schapira K, Roth M (1969). The Relationship between Premature Death and Affective Disorders. British Journal of Psychiatry, 115: 1277 - 82.
54046	Kessler RC, Berglund P, Borges G, et al (2005). Trends in suicide ideation, plans, gestures, and attempts in the United States, 1990-1992 to 2001-2003. JAMA, 293(20): 2487-95.
54042	Kessler RC, Berglund PA, Borges G, et al (2007). Smoking and suicidal behaviors in the National Comorbidity Survey: replication. J Nerv Ment Dis, 195(5): 369-77.
52679	Kessler RC, McRae JA (1983). Trends in the relationship between sex and attempted suicide. J Health Sci Behav, 24: 98-110.
6745	Kessler RC, Sonnega A, Bromet E, et al (1995). Posttraumatic stress disorder in the national comorbidity survey. Arch Gen Psychiatry, 52(12): 1048-60.
52671	Kevan SM (1980). Perspectives on season of suicide: a review. Soc Sci Med, 14(4): 369-78.
47072	Khalsa HM, Salvatore P, Hennen J, et al (2008). Suicidal events and accidents in 216 first-episode bipolar I disorder patients: predictive factors. Journal of Affective Disorders, 106: 179-84.
54009	Kikuchi N, Ohmori-Matsuda K, Shimazu T, et al (2009). Pain and risk of completed suicide in Japanese men: a population-based cohort study in Japan (Ohsaki Cohort Study). J Pain Symptom Management, 37(3): 316-24.
52451	Kim HS, Kim HS (2008). Risk factors for suicide attempts among Korean adolescents. Child Psychiatry Hum Dev, 39: 221-35.
84118	Kim MS, Hong YC, Yook JO, et al (2017). Effects of perceived job insecurity on Effects of perceived job insecurity on depression, suicide ideation, and decline in self-rated health in Korea: a population-based panel study., 90: 663-71.
49903	Kim YS, Koh YJ, Leventhal B (2005). School bullying and suicidal risk in Korean middle school students. Pediatrics, 115(2): 357-63.
52148	Kim YS, Leventhal B (2008). Bullying and suicide. A review. Int J Adolesc Med Health, 20(2): 133-54.
54280	Kim YS, Leventhal BL, Koh YJ, Boyce WT (2009). Bullying increases suicide risk: prospective study of Korean adolescents. Arch Suicide Res, 13(1): 15-30.
11036	Kimerling R, Calhoun KS (1994). Somatic symptoms, social support and treatment seeking among sexual assault victims. Journal of Consulting and Clinical Psychology, 62(2): 333-40.
77478	Kimerling R, Makin-Byrd K, Louzon S, et al (2015). Military sexual trauma and suicide mortality. Am J Prev Med, (Epub ahead of print).
32364	King DW, King LA, Gudanowski DM, et al (1995). Alternative representations of war zone stressors: relationships to posttraumatic stress disorder in male and female Vietnam veterans. Journal of Abnormal Psychology, 104(1): 184-96.

49902	King M, Semlyen J, Tai SS, et al (2008). A systemic review of mental disorder, suicide, and deliberate self-harm in lesbian, gay and bisexual
	people. BMC Psychiatry, 8: 70.
78019	Kitchener S, Nasveld P, Edstein MD (2007). Short report: tafenoquine for the treatment of recurrent plasmodium vivax malaria. Am J Trop Med Hyg, 76(3): 494-6.
52660	Klerman GL (1987). Clinical epidemiology of suicide. J Clin Psychiatry, 48(12 (Suppl)): 33-8.
79750	Kline A, Weiner MD, Interian A, et al (2016). Morbid thoughts and suicidal ideation in Iraq war veterans: the role of direct and in direct killing in combat. Depress Anxiety, 33: 473-82.
53680	Klomek AB, Marrocco F, Kleinman M, et al (2007). Bullying, depression, and suicidality in adolescents. J Am Acad Child Adolesc Psychiatry, 46(1): 40-9.
53677	Klomek AB, Sourander A, Kumpulainen K, et al (2008). Childhood bullying as a risk for later depression and suicidal ideation among Finnish males. J Affect Disord, 109: 47-55.
53679	Klomek AB, Sourander A, Niemela S, et al (2009). Childhood bullying behaviors as a risk for suicide attempts and completed suicides: a population-based birth cohort study. J Am Acad Child Adolesc Psychiatry, 48(3): 254-61.
68400	Knapik JJ, Marin RE, Grier TL et al (2009). A systematic review of post- deployment injury-related mortality among military personnel deployed to conflict zones. BMC Public Health, 9: 231.
79694	Knox KL (2008). Epidemiology of the relationship between traumatic experience and suicidal behaviors. PTSD Research Quarterly, 19(4): 1-8.
83480	Kochanski-Ruscio K, Nademin E, Perera K, et al (2016). An examination of United States Air Force suicide decendents based on documented suicide attempt histories. Arch Suicide Res, 1-12.
77854	KokoAung E, Aromataris E (2012). The association between suicidality and treatment with selective serotonin reuptake inhibitors in older people with major depression: a systematic review. The JBI database of systematic review and implementation reports, 10(56).
4209	Korn ML, Kotler M, Molcho A, et al (1992). Suicide and violence associated with panic attacks. Biol Psychiatry, 31: 607-12.
77764	Kosidou K, Dalman C, Fredlund P, et al (2014). School performance and the risk of suicide attempts in young adults: a longitudinal population-based study. Psychol Med, 44: 1235-43.
53988	Koskinen O, Pukkila K, Hakko H, et al (2002). Is occupation relevant in suicide? J Affect Disord, 70: 197-203.
53461	Kovacic Z, Henigsberg N, Pivac N, et al (2008). Platelet serotonin concentration and suicidal behaviour in combat related posttraumatic stress disorder. Prog Neuropsychopharmacol Biol Psychiatry, 32: 544-51.
1757	Kramer TL, Lindy JD, Green BL, et al (1994). The comorbidity of post-traumatic stress disorder and suicidality in Vietnam veterans. Suicide Life Threat Behav, 24(1): 58-67.
77366	Kraus JE, Horrigan JP, Carpenter DJ, et al (2010). Clinical features of patients with treatment-emergent suicidal behavior following initiation of paroxetine therapy. J Affect Disord, 120(1): 40-7.
77830	Krentz AJ, Fujioka K, Hompesch M (2016). Evolution of pharmacological obesity treatments: focus on adverse side-effects profiles. Diabetes Obes Metab, [Epub ahead of print]: doi:10.111/dom.12657.
45493	Krysinska K, Heller TS, De Leo D (2006). Suicide and deliberate self- harm in personality disorders. Curr Opin in Psychiatry, 19: 95-101.
54368	Kuehn BM (2008). FDA warns of adverse events linked to smoking cessation drug and antiepileptics. JAMA, 299(10): 1121-2.

53448	Kuehn BM (2009). Soldier suicide rates continue to rise. Military, scientists work to stem the tide. JAMA, 301(11): 111, 113.
54369	Kuehn BM (2009). Studies linking smoking-cessation drug with suicide risk spark concerns. JAMA, 301(10): 1007-8.
54726	Kuehn JK, Kuehn JL (1975). Conspiracy of silence: psychiatric counselling with students at high risk for academic failure. Am J Psychiatry, 132(11): 1207-9.
11037	Kushner MG, Riggs DS, Foa EB, et al (1992). Perceived controllability and the development of posttraumatic stress disorder (PTSD) in crime victims. Behav Res Ther, 31(1): 105-10.
85928	LaFauci Schutt JM, Marotta SA (2011). Personal and environmental predictors of posttraumatic stress in emergency management professionals. Psychological Trauma, 3(1): 8-12.
28601	Lambert MT, Fowler DR (1997). Suicide risk factors among veterans: risk management in the changing culture of the Department of Veterans Affairs. The Journal of Behavioral Health Sciences & Research, 24(3): 350-58.
54017	Lapierre YD (2003). Suicidality with selective serotonin reuptake inhibitors: valid claim? J Psychiatry Neurosci, 28(5): 340-7.
4198	LaRosa JC (1995). Cholesterol lowering and morbidity and mortality. Current Opinion in Lipidology, 6: 62-65.
52446	Laser J, Luster T, Oshio T (2007). Risk and promotive factors related to depressive symptoms among Japanese youth. Am J Orthopsychiatry, 77(4): 523-33.
83765	Law YW, Yip PSF, Zhang Y, et al (2014). The chronic impact of work on suicides and under-utilization of psychiatric and psychosocial services. J Affect Disord, 168: 254-61.
79645	LeardMann CA, Powell TM, Smith TC, et al (2013). [Erratum] Risk factors associated with suicide in current and former US military personnel. JAMA, 310(10): 1076. ID: 0.
79644	LeardMann CA, Powell TM, Smith TC, et al (2013). Risk factors associated with suicide in current and former US military personnel. JAMA, 310(5): 496-506.
78017	Leary KJ, Riel MA, Roy MJ, et al (2009). A randomized, double-blind, safety and tolerability study to assess the ophthalmic and renal effects of tafenoquine 200 mg weekly versus placebo for 6 months in healthy volunteers. Am J Trop Med Hyg, 81(2): 356-62.
76984	Lebain P, Juliard C, Davy JP, et al (2000). Neuropsychiatric symptoms in preventive antimalarial treatment with mefloquine: apropos of 2 cases. Encephale, 26(4): 67-70. [Abstract]
79748	LeBouthillier DM, McMillan KA, Thibodeu MA, et al (2015). Types and number of traumas associated with suicidal ideation and suicide attempts in PTSD: findings from a U.S. nationally representative sample. Journal of Traumatic Stress, 28: 183-90.
53998	Lee WJ, Alavanja MCR, Hoppin JA, et al (2007). Mortality among pesticide applicators exposed to Chlorpyrifos in the Agricultural Health Study. Environ Health Perspect, 115(4): 528-34.
78022	Lell B, Faucher J-F, Missinou MA, et al (2000). Malaria chemoprophylaxis with tafenoquine: a randomised study. Lancet, 355(9220): 2041-5.
54003	Lepine J-P, Briley M (2004). The epidemiology of pain in depression. Human Psychopharmacology, 19: S3-7.
4202	Lepine JP, Chignon JM, Teherani M (1993). Suicide attempts in patients with panic disorder. Arch Gen Psychiatry, 50: 144-9.
53675	Lester D (1973). [Comment] Suicide in released prisoners of war. JAMA, 225(7): 747.
4204	Lester D (1994). Experience of loss and subsequent suicide. Perceptual and Motor Skills, 79: 730.

54764	Lester D (1997). The role of shame in suicide. Suicide & Life-Threatening Behaviour, 27(4): 352-61.
54686	Lester D (1998). The association of shame and guilt with suicidality. The Journal of Social Psychology, 138(4): 535-6.
54282	Lester D (2002). Serum cholesterol levels and suicide: a meta-analysis. Suicide & Life-Threatening Behaviour, 32(3): 333-46.
54283	Lester D (2003). [Comment] Serum cholesterol levels and suicide: A further comment. Suicide & Life-Threatening Behaviour, 33(3): 338-9. Comment on ID: 54282.
52685	Lester D (2006). Suicide and Islam. Arch Suicide Res, 10(1): 77-97.
4203	Lester D, Moderski T (1995). The Imposter Phenomenon in Adolescents. Psychological Reports, 76: 466.
77474	Levi-Belz Y, Zerach G, Solomon Z (2015). Suicide ideation and deliberate self-harm among ex-prisoners of war. Arch Suicide Res, 19(2): 231-48.
39368	Li J, Hansen Precht D, Mortensen PB, et al (2003). Mortality in parents after death of a child in Denmark: a nationwide follow-up study. Lancet, 361: 363-7.
77377	Li J, Vestergaard M, Cnattingius S, et al (2014). Mortality after parental death in childhood: a nationwide cohort study from three Nordic countries. PLoS Med, 11(7): e1001679.
54001	Lima CS, Ribeiro-Carvalho A, Filgueiras CC, et al (2009). Exposure to methamidophos at adulthood elicits depressive-like behavior in mice. NeuroToxicology, 30: 471-8.
54035	Limosin F, Loze J-Y, Philippe A, et al (2007). Ten-year prospective follow-up study of the mortality by suicide in schizophrenic patients. Schizophrenia Research, 94: 23-8.
45576	Links PS, Gould B, Ratnayake R (2003). Assessing suicidal youth with antisocial, borderline, or narcissistic personality disorder. Can J Psychiatry, 48(5): 301-10.
85929	Liu B, Tarrigan LH, Bromet EJ, et al (2014). World trade centre disaster exposure-related probable posttraumatic stress disorder among responders and civilians: A meta-analysis. PLoS One, 9(7): e101491.
79780	Liu RT, Miller I (2014). Life events and suicidal ideation and behavior: a systematic review. Clin Psychol Rev, 34: 181-92.
52440	Liu X, Li H, Qin W, et al (2006). Association of TPH1 with suicidal behaviour and psychiatric disorders in the Chinese population. J Med Genet, 43: e4.
78014	Llanos-Cuentas A, Lacerda MV, Rueangweerayut R, et al (2014). Tafenoquine plus chloroquine for the treatment and relapse prevention of plasmodium vivax malaria (DETECTIVE): a multicentre, double-blind, randomised, phase 2b dose-selection study. Lancet, 383(9922): 1049-58.
4190	Lloyd GG (1993). Acute behaviour disturbances. Journal of Neurology. Neurosurgery, and Psychiatry, 56: 1149-56.
76985	Lobel HO, Coyne PE, Rosenthal PJ (1998). Drug overdoses with antimalarial agents: prescribing and dispensing errors. JAMA, 280(17): 1483.
54724	Loebel JP, Loebel JS, Dager SR, et al (1991). Anticipation of nursing home placement may be a precipitant of suicide among the elderly. JAGS, 39: 407-8.
54032	Loftis JM, Matthews AM, Hauser P (2006). Psychiatric and substance use disorders in individuals with hepatitis C. Epidemiology and management. Drugs, 66(2): 155-74.
77485	Logan BK, Yeakel JK, Goldfogel G, et al (2012). Dextromethorphan abuse leading to assault suicide, or homicide. J Forensic Sci, 57(5): 1388-94.

Logan J, Bohnert A, Spies E, et al (2013). Suicidal ideation among young Afghanistan/Iraq war veterans and civilians: Individual, social, and environmental risk factors and perception of unmet mental healthcare needs. Psychiatry Res, [Epub ahead of print].
London L, Flisher AJ, Wessling C, et al (2005). Suicide and exposure to organophosphate insecticide: cause or effect? Am J Ind Med, 47: 308-21.
Lucaclu LA, Dumitrascu DL (2015). Depression and suicide ideation in chronic hepatitis C patients untreated and treated with interferon: prevalence, prevention, and treatment. Ann Gastroenterol, 28(4): 440-7.
Ludot M, Mouchabac S, Ferreri F (2015). Inter-relationships between isotretinoin treatment and psychiatric disorders: depression, bipolar disorder, anxiety, psychosis and suicide risks. World J Psychiatr, 5(2): 222-7.
Macfarlane GJ, McBeth J, Silman AJ (2001). Widespread body pain and mortality: prospective population based study. BMJ, 323: 1-4.
Mackelprang JL, Bombardier CH, Fann JR (2014). Rates and predictors of suicidal ideation during the first year after traumatic brain injury. Am J Public Health, 104(7): e100-7.
MacLeod AK, Pankhania B, Lee M, et al (1997). Parasuicide, depression and the anticipation of positive and negative future experiences. Psychol Med, 27: 973-7.
Magin P, Pond D, Smith W (2005). Isotretinoin, depression and suicide: a review of the evidence. British Journal of General Practice, 55: 134-8.
Mahon MJ, Tobin JP, Cusack DA, et al (2005). Suicide among regular- duty military personnel: a retrospective case-control study of occupation- specific risk factors for workplace suicide. Am J Psychiatry, 162: 1688-96.
Mahuen S, Metzler TJ, Bosch J, et al (2012). Killing in combat may be independently associated with suicidal ideation. Depress Anxiety, 29(11): 918-23.
Manchikanti L, Helm II S, Fellows B, et al (2012). Opioid epidemic in the United States. Pain Physician, 15: ES9-38.
Manguno-Mire G, Sautter F, Lyons J, et al (2007). Psychological distress and burden among female partners of combat veterans with PTSD. J Nerv Ment Dis, 195(2): 144-51.
Mann J, McBride A, Brown RP, et al (1992). Relationship between central and peripheral serotonin indexes in depressed and suicidal psychiatric inpatients. Arch Gen Psychiatry, 49: 442-6.
Mann JJ (2003). Neurobiology of suicidal behaviour. Nat Rev Neurosci, 4(10): 819-28.
Mann JJ, Currier D (2007). A review of prospective studies of biologic predictors of suicidal behavior in mood disorders. Arch Suicide Res, 11(1): 3-16.
March J, Sareen J, Gawaziuk JP, et al (2014). Increased suicidal activity following major trauma: a population-based study. J Trauma Acute Care Surg, 76: 180-4.
Maris RW (1997). Social and familial risk factors in suicidal behavior. The Psychiatric Clinics of North America, 20(3): 519-50.
Martikainen P, Valkonen T (1996). Mortality after the death of a spouse: rates and causes of death in a large Finnish cohort. American Journal of Public Health, 86(8): 1087-93.
Martin RL, Cloninger CR, Guze SB, et al (1985). Mortality in a follow-up of 500 psychiatric outpatients: II. cause specific mortality. Arch Gen Psychiatry, 42: 58-66.
Masi G, Tomaiuolo F, Sbrana B, et al (2001). Depressive symptoms and academic self-image in adolescence. Psychopathology, 34: 57-61.

T
Massion AO, Warshaw MG, Keller MB (1993). Quality of life and
psychiatric morbidity in panic disorder and generalised anxiety disorder.
Am J Psychiatry, 150(4): 600-7.
Mather AA, Cox BJ, Enns MW, et al (2009). Associations of obesity with
psychiatric disorders and suicidal behaviors in a nationally representative
sample. J Psychosom Res, 66: 277-85.
Matsumoto Y, Shimizu K, Kinoshita H, et al (2010). Suicide associated
with corticosteroid use during chemotherapy: case report. Jpn J Clin
Oncol, 40(2): 174-6.  Mattunen MJ, Aro HM, Henriksson MM, et al (1991). Mental disorders in
adolescent suicide: DSM III R axes I and II diagnoses in suicides among
13 to 19 year olds in Finland. Arch Gen Psychiatry, 48: 834-9.
Maxwell NM, Nevin RL, Stahl S, et al (2015). Prolonged neuropsychiatric
effects following management of chloroquine intoxication with
psychotropic polypharmacy. Clin Case Rep, 3(6): 379-387.
McCarthy JF, Valenstein M, Kim HM, et al (2009). Suicide mortality
among patients receiving care in the veterans health administration
health system. Am J Epidemiol, 169(8): 1033-8.
McCarthy S (2015). Malaria prevention, mefloquine neurotoxicity,
neuropsychiatric illness, and risk-benefit analysis in the Australian
Defence Force. Journal of Parasitology Research, ID287651: 23 pages.
McCrea M, Guskiewicz K, Randolph C, et al (2013). Incidence, clinical
course, and predictors of prolonged recovery time following sport-related
concussion in high school and college athletes. J Int Neuropsychol Soc,
19(1): 22-33.
McDermott E, Roen K, Scourfield J (2008). Avoiding shame: young LGBT
people, homophobia and self-destructive behaviours. Culture, Health &
Sexuality, 10(8): 815-29.
McFarlane AC, Hodson SE, Van Hooff M, et al (2011). Mental health in
the Australian Defence Force: full report. 2010 ADF Mental Health and
Wellbeing Study. Department of Defence, Canberra.
McKenzie D (2010). An holistic view of post-traumatic stress disorder.
Journal of Military Health, 18: 24-6.
McLaughlin KA, Koenen KC, Friedman MJ, et al (2015). Sub-threshold
post traumatic stress disorder in the WHO world mental health surveys.
Biol Psychiatry, 77(4): 375-84.
Meduri GU, Shih M-C (2012). [Comment] Acute clinical deterioration with
increased systemic inflammation as a risk factor for depression: an
alternative interpretation of the data. Am J Psychiatry, 169(8): 867.  Mehlum L, Friis S, Karterud S (1994). The longitudinal pattern of suicidal
behaviour in borderline personality disorder: a prospective follow-up
study. Acta Psychiatr Scand, 90: 124-30.
Meltzer HY, Alphs L, Green AI, et al (2003). Clozapine treatment for
suicidality in schizophrenia. Arch Gen Psychiatry, 60: 82-91.
Mezuk B, Lohman M, Leslie M, et al (2015). Suicide risk in nursing
homes and assisted living facilities: 2003-2011. Am J Public Health,
105(7): 1495-1502.
Micciolo R, Williams P, Zimmermann-Tansella CH, et al (1991).
Geographical and urban-rural variation in the seasonality of suicide:
some further evidence. J Affect Disord, 21: 39-43.
Miller BC, Monson BH, Norton MC (1995). The effects of forced sexual
intercourse on white female adolescents. Child Abuse & Neglect, 19(10):
1289-01.
Miller M, Barber C, Azrael D, et al (2009). Suicide among US veterans: a
prospective study of 500,000 middle-aged and elderly men. Am J
Epidemiol, 170(4): 494-500.

83210	Milner A, Currier D, LaMontagne AD, et al (2017). Psychosocial job stressors and thoughts about suicide among males: a cross-sectional
83510	study from the first wave of the Ten to Men cohort. Public Health.  Milner A, Page K, Witt K, et al (2016). Psychosocial working conditions and suicide ideation: Evidence from a cross-sectional survey of working Australians. J Occup Environ Med, 58(6): 584.
83509	Milner A, Spittal MJ, Pirkis J, et al (2017). Low control and high demands at work as risk factors for suicide: An Australian national population-level case-control study. Psychosom Med, 79(3): 358-364.
83723	Milner A, Witt K, LaMontagne AD, et al (2017). Psychosocial job stressors and suicidality: a meta-analysis and systematic review. Occup Environ Med, 75(4): 245-53.
52438	Misfeldt J, Senderovitz F (1989). Suicide in Greenland. Arctic Med Res, 48: 122-3.
86476	Misra M, Greenberg N, Hutchinson C, et al (2009). Psychological impact upon London ambulance service of the 2005 bombings. Occup Med, 59(6): 428-33.
77932	Mitchell JE, Crosby R, de Zwaan M, et al (2013). Possible risk factors for increased suicide following bariatric surgery. Obesity, 21(4): 665-72.
4185	Modestin J, Villiger C (1989). Follow-up study on borderline versus nonborderline personality disorders. Comprehensive Psychiatry, 30(3): 236-44.
77484	Modi D, Bhalavat R, Patterson JC (2013). Suicidal and homicidal behaviors related to dextromethorphan abuse in a middle-aged woman. J Addict Med, 7(2): 143-4.
78066	Mohan D, Mohandas E, Rajat R, et al (1981). Chloroquine psychosis: a chemical psychosis. J Nat Med Assoc, 73(11): 1073-6.
54765	Mokros HB (1995). Suicide and shame. The American Behavioral Scientist, 38(8): 1091-103.
79749	Monteith LL, Bahraini NH, Matarazzo BB, et al (2016). The influence of gender on suicidal ideation following military sexual trauma among Veterans in the Veterans Health Administration. Psychiatry Res, 244: 257-65.
77797	Moore TJ, Furberg CD, Glenmullen J, et al (2011). Suicidal behavior and depression in smoking cessation treatments. PLoS One, 4(11): e27016.
4208	Morano CD, Cisler RA, Lemerond J (1993). Risk factors for adolescent suicidal behaviour: Loss, insufficient familial support, and hopelessness. Adolescence, 28(112): 851-65.
77792	Moreland CS, Bonin L (2015). Effect of antidepressants on suicide risk in children and adolescents. Retrieved 17 March 2016, from www.uptodate.com/contents/ effect-of-antidepressants-on-suicide-risk-in-children-and-adolescents
86129	Morgan PM (2016). The psychological impact of mass casualty incidents on first responders: A systematic review. Journal of Emergency Management, 14: 213-26.
22686	Morris P, Raphael B, Bordujenko A (eds.) (1999). Repatriation Medical Authority Consensus Conference Proceedings: Stress and Challenge - Health and Disease, Brisbane February 9-11, 1998. Repatriation Medical Authority, Brisbane.
54690	Mortensen PB (1999). [Comment] Can suicide research lead to suicide prevention? Acta Psychiatr Scand, 99: 397-8. Comment on ID: 54689.
52669	Moscicki EK (1989). Epidemiologic surveys as tools for studying suicidal behavior: A review. Suicide & Life-Threatening Behaviour, 191(1): 131-46.
1761	Moscicki EK (1995). Epidemiology of suicidal behaviour. Suicide and Life-Threatening Behaviour, 25(1): 22-35.

54272	Moscicki EK (2004). [Comment] Opportunities of life. Preventing suicide in elderly patients. Arch Intern Med, 164: 1171-2. Comment on ID: 54271.
79696	Mota NP, Tsai J, Sareen J, et al (2016). High burden of subthreshold DSM-5 post-traumatic stress disorder in U.S. military veterans. World Psychiatry, 15(2): 185-6.
72426	Mott JM, Graham DP, Teng EJ (2012). Perceived threat during deployment: risk factors and relation to Axis I disorders. Psychological Trauma: Theory, Research Practice and Policy, 4(6): 587-595.
77471	Mukamal KJ, Rimm EB, Kawachi I, et al (2010). Body mass index and risk of suicide among one million US adults. Epidemiol, 21(1): 82-6.
77481	Mula M, Sander JW (2015). Suicide and epilepsy: do antiepileptic drugs increase the risk? Expert Opin Drug Saf, 14(4): 553-8.
4195	Muldoon MF, Rossouw JE, Manuck SB, et al (1993). Low or lowered cholesterol and risk of death from suicide and trauma. Metabolism, 42(9 S 1): 45-56.
11039	Murdoch M, Nichol KL (1995). Women veterans' experiences with domestic violence and with sexual harassment while in the military. Arch Fam Med, 4(5): 411-8.
52672	Murphy GE (1983). Problems in studying suicide. Psychiatric Developments, 4: 339-50.
1762	Murphy GE, Wetzel RD (1990). The lifetime risk of suicide in alcoholism. Arch Gen Psychiatry, V47: 383-92.
53456	Nad S, Marcinko D, Vuksan-Eusa B, et al (2008). Spiritual well-being, intrinsic religiosity, and suicidal behavior in predominantly Catholic Croatian war veterans with chronic posttraumatic stress disorder. A case control study. J Nerv Ment Dis, 196(1): 79-83.
54290	Naranjo CA, Shear NH, Lanctot KL (1992). Advances in the diagnosis of adverse drug reactions. J Clin Pharmacol, 32: 897-904.
78015	Nasveld P, Kitchener S, Edstein M, et al (2002). Comparison of tafenoquine (WR238605) and primaquine in the post-exposure (terminal) prophylaxis of vivax malaria in Australian Defence Force personnel. Trans R Soc Trop Med Hyg, 96(6): 683-4.
77907	Nasveld PE, Edstein MD, Reid M, et al (2010). Randomized, double-blind study of the safety, tolerability, and efficacy of tafenoquine versus mefloquine for malaria prophylaxis in nonimmune subjects. Antimicrob Agents Chemother, 54(2): 792-8.
9596	Nefzger MD (1970). Follow-up studies of World War II and Korean War Prisoners. I Study Plan and Mortality findings. Am J Epidemiol, 91(2): 123-38.
79697	Nelson H, Denneson L, Low A, et al (2015). Systematic review of suicide prevention in veterans. Evidence-based Synthesis Program (ESP) Center Portland VA Medical Center. Dept of Veterans Affairs.
75199	Nevin RL (2012). [Comment] Hallucinations and persecutory delusions in Mefloquine-Associated suicide. Am J Forensic Med Pathol, 33(2): e8.
75201	Nevin RL (2012). Limbic encephalopathy and central vestibulopathy caused by mefloquine: A case report. Travel Medicine and Infectious Disease, 10: 144-51.
76942	Nevin RL (2015). Mefloquine and posttraumatic stress disorder. Textbook of military medicine: Forensic and Ethical Issues in Military Behavioural Health, 19: 277-96. Borden Institute, Washington D.C.
60085	Nevin RL (2016). The mefloquine intoxication syndrome: a significant potential confounder in the diagnosis and management of PTSD and other chronic deployment-related neuropsychiatric disorders. Post-Traumatic Stress Disorder and Related Diseases in Combat Veterans, : Springer International Switzerland.

79646	Nevin RL, Ritchie EC, Bryan CJ, et al (2013). [Comments] Suicides among military personnel. JAMA, 310(23): 2563-6. Comments on ID: 79644.
73303	Nevoralová Z, Dvoráková D (2013). Mood changes, depression and suicide risk during isotretinoin treatment: a prospective study. Int J Dermatol, 52(2): 163-8.
77477	Newton-John TR (2014). Negotiating the Maze: risk factors for suicidal behavior in chronic pain patients. Curr Pain Headache Rep, 18(9): 447.
77721	Niedzwiedz C, Haw C, Hawton K, et al (2014). The definition and epidemiology of clusters of suicidal behavior: a systematic review. Suicide & Life-Threatening Behaviour, 44(5): 569-81.
1763	Nielson B, Wang GA, Bille-Brahe U (1990). Attempted suicide in Denmark. Five years follow-up. Acta Psychiatr Scand, V81: 250-4.
81521	Nock M, Deming C, Fullerton C, et al (2013). Suicide among soldiers: A review of psychosocial risk and protective factors. Psychiatry, 76(2): 97-125.
52449	Nock MK, Borges G, Bromet EJ, et al (2008). Cross-national prevalence and risk factors for suicidal ideation, plans and attempts. Br J Psychiatry, 192: 98-105.
77378	Nock MK, Green JG, Hwang I, et al (2013). Prevalence, correlates and treatment of lifetime suicidal behavior among adolescents: results from the national comorbidity survey replication - adolescent supplement (NCSA). JAMA Psychiatry, 70(3).
54454	Nock MK, Hwang I, Sampson N, et al (2009). Cross-national analysis of the associations among mental disorders and suicidal behaviour: findings from the WHO World Mental Health surveys. PLoS Med, 6(8): e1000123.
55084	Nock MK, Kessler RC (2006). Prevalence of and risk factors for suicide attempts versus suicide gestures: analysis of the National Comorbidity Survey. Journal of Abnormal Psychology, 115(3): 616-23.
1764	Nordentoft M, Rubin P (1993). Mental illness and social integration among suicide attempters in Copenhagen: comparison with the normal population and a four-year follow-up study of 100 patients. Acta Psychiatr Scand, 88: 278-85.
47151	Norris FH (1990). Screening for traumatic stress: a scale for use in the general population. Journal of Applied Social Psychology, 20(20): 1704-18.
86477	North CS, Tivis L, McMillen JC, et al (2002). Coping, functioning, and adjustment of rescue workers after the Oklahoma City bombing. J Trauma Stress, 15(3): 171-5.
1765	Noyes R Jr (1991). Suicide and panic disorder: a review. J Affective Disorders, 22: 1-11.
1766	Noyes R, Christiansen J, Clancy J, et al (1991). Predictors of serious suicide attempts among patients with panic disorder. Comprehensive Psychiatry, 32(3): 261-7.
77946	Nugent KL, Paksarian D, Mojtabai R (2011). Non-affective acute psychoses: Uncertainties on the way to DSM-V and ICD-11. Curr Psychiatry Rep, 13(3): 203-10.
75635	O'Brien BS, Sher L (2013). Military sexual trauma as a determinant in the development of mental and physical illness in male and female veterans. Int J Adolesc Med Health, 25(3): 269-74.
45596	O'Connell KA, Wilkin JK, Pitts M (2002). [Comment] Isotretinoin (Accutane) and serious psychiatric adverse events. J Am Acad Dermatol, 48(2): 306-8.
54020	O'Donnell J (2004). Polar hysteria: an expression of hypervitaminosis A. Am J Ther, 11: 507-16.
	<u> </u>

54900	Oedegaard KJ, Angst J, Neckelmann D, et al (2005). Migraine aura without headache compared to migraine with aura in patients with
52969	affective disorders. J Headache Pain, 6: 378-86.  Ohgami H, Terao T, Shiotsuki I, et al (2009). Lithium levels in drinking water and risk of suicide. The British Journal of Psychiatry, 194(5): 464-5.
81436	Olenick M, Flowers M, Diaz VJ (2015). US veterans and their unique issues: enhancing health care professional awareness. Adv Med Educ Pract, 6: 635-9.
86456	Omerov P, Pettersen R, Titelman D, et al (2016). Encountering the body at the site of the suicide: A population-based survey in Sweden. Suicide and Life-Threatening Behavior, 47(1): 38-47.
54827	Orbach I (2008). Mental pain, pain-producing constructs, the suicide body, and suicide. Relating to Self-harm and Suicide: Psychoanalytic Perspectives on Practice, Theory and Prevention, Chapter 7: 80-92. Routledge, New York.
52445	O'Reilly D, Rosato M, Connolly S, et al (2008). Area factors and suicide: 5-year follow-up of the Northern Ireland population. Br J Psychiatry, 192: 106-11.
52153	O'Shea B (2000). Self-harm and unemployment. Hospital Medicine, 61(7): 195-8.
52905	Ostry A, Maggi S, Tansey J, et al (2007). The impact of psychosocial work conditions on attempted and completed suicide among western Canadian sawmill workers. Scandinavian Journal of Public Health, 35: 265-71.
28653	O'Toole BI, Cantor C (1995). Suicide risk factors among Australian Vietnam era draftees. Suicide & Life-Threatening Behaviour, 25(4): 475-88.
4191	Owens D (1994). Cholesterol and violent death: Clinical importance is questionable. British Medical Journal, 309(6963): 1228.
54737	Owens D, Horrocks J, House A (2002). Fatal and non-fatal repetition of self-harm. Br J Psychiatry, 181: 193-9.
53987	Page AN, Fragar LJ (2002). Suicide in Australian farming, 1988-1997. Aust N Z J Psychiatry, 36: 81-5.
21880	Page WF, Tanner CM (2000). [Comment] Parkinson's disease and motor-neuron disease in former prisoners-of-war. The Lancet, 355(9206): 843. Comment on ID: 21879.
1767	Pandey GN, Pandey SC, Dwivedi Y, et al (1995). Platelet serotonin-2A receptors: a potential biological marker for suicidal behaviour. Am J Psychiatry, 152: 850-5.
54289	Paolucci EO, Genuis ML, Violato C (2001). A meta-analysis of the published research on the effects of child sexual abuse. J Psychol, 135(1): 17-36.
54040	Pariente A, Daveluy A, Laribiere-Benard A, et al (2009). Effect of date of drug marketing on disproportionality measures in pharmacovigilance. Drug Safety, 32(5): 441-7.
4177	Paris J, Nowlis D, Brown R (1989). Predictors of suicide in borderline personality disorder. Can J Psychiatry, 34: 8-9.
76941	Park H, Reber BH, Chon M-G (2015). Tweeting as health communication: health organizations' use of twitter for health promotion and public engagement. Journal of Health Communication, [Epub ahead of print]0: 1-11.
77724	Park S-J, Lee HB, Ahn MH, et al (2015). Identifying clinical correlates for suicide among epilepsy patients in South Korea: a case-control study. Epilepsia, 56(12): 1966-72.
83722	Peltzer K, Yi S, Pengpid S (2017). Suicidal behaviors and associated factors among university students in six countries in the Association of Southeast Asian Nations (ASEAN). Asian J Psychiatr, 26: 32-8.

	Pereira A, Gitlin MJ, Gross RA, et al (2013). Suicidality associated with
77561	anti-epileptic drugs: implications for the treatment of neuropathic pain and
	fibromyalgia. Pain, 154(3): 345-9.
77562	Perera S, Eisen R, Bawor M, et al (2015). Association between body
	mass index and suicidal behaviors: a systematic review protocol.
	Systematic Reviews, 4: 52.
54570	Perez-Olmos I, Rodriguez-Sandoval E, et al (2007). Psychiatric and
	social characterisation of suicide attempts treated at a children's clinic,
	2003-2005. Revista De Salud Publica, 9(2): 230-40.
50074	Peter T, Roberts LW, Buzdugan R (2008). Suicidal ideation among
53674	Canadian youth: a multivariate analysis. Archives of Suicide Research,
	12(3): 263-75.
77007	Peterhansel C, Petroff D, Zlinitzke G, et al (2013). Risk of completed
77927	suicide after bariatric surgery: a systematic review. Obesity Reviews,
	14(5): 369-82.
4=00	Petronis KR, Samuels JF, Moscicki EK, et al (1990). An epidemiological
1768	investigation of potential risk factors for suicide attempts. Soc Psychiatry
	Psychiatr Epidemiol, 25: 193-9.
50400	Pfeffer CR, Lipkins R, Plutchik R, et al (1988). Normal children at risk for
52163	suicidal behaviour: a two-year follow-up study. J Am Acad Child Adolesc
	Psychiatry, 27(1): 34-41.
53986	Pickett W, King WD, Lees REM, et al (1998). Suicide mortality and
	pesticide use among Canadian farmers. Am J Ind Med, 34: 364-72.
00400	Pietrzak RH, Feder A, Singh R, et al (2014). Trajectories of PTSD risk
86468	and resilience in World Trade Center responders: an 8-year prospective
	cohort study. Psychol Med, 44(1): 205-19.
	Pietrzak RH, Schechter CB, Bromet EJ, et al (2012). The burden of full
70933	and subsyndromal posttraumatic stress disorder among police involved in
	the World Trade Center rescue and recovery effort. J Psychiatr Res,
	46(7): 835-42.
77809	Pigeon WR, Pinquart M, Connor K (2012). Meta-analysis of sleep disturbance and suicidal thoughts and behaviors. J Clin Psychiatry, 73(9):
11009	e1160-7.
	Pinna F, Sanna L, Perra V, et al (2014). Long-term outcome of
77600	schizoaffective disorder. Are there any differences with respect to
77000	
	schizophrenia. Riv Psichiatr, 49(1): 41-9.  Pirkis J (2004). [Comment] Borderline personality disorder, drug use
	disorder, and worsening depression or substance abuse are significant
49801	predictors of suicide attempts in people with Axis I and II disorders. Evid
	Based Ment Health, 7(1): 25. Comment on ID: 49800.
	Platt S (1987). The aftermath of Angie's overdose: is soap (opera)
52164	damaging to your health? BMJ, 294: 954-7.
	Pompili M, Di Cosimo D, Innamorati M, et al (2009). Psychiatric
	comorbidity in patients with chronic daily headache and migraine: a
54771	selective overview including personality traits and suicide risk. J
	Headache Pain, 10: 283-90.
	Pompili M, Girardi P, Tatarelli G, et al (2006). Suicide and attempted
54288	suicide in eating disorders, obesity and weight-image concern. Eating
	Behaviors, 7: 384-94.
	Pompili M, Girardi P, Tatarelli R (2006). Death from suicide versus
54287	mortality from epilepsy in the epilepsies: A meta-analysis. Epilepsy &
	Behavior, 9: 641-8.
	Pompili M, Innamorati M, Szanto K, et al (2011). Life events as
79782	precipitants of suicide attempts among first-time suicide attempters,
13102	repeaters, and non-attempters. Psychiatry Res, 186: 300-5.
	Pompili M, Mancinelli I, Girardi P, et al (2004). Suicide in anorexia
54286	nervosa: a meta-analysis. Int J Eat Disord, 36: 99-103.
	indivosa. a meta-analysis. Int o Lat Disord, 50. 33-105.

<u> </u>	
	Preda A (2008). Suicidality and depression: medicolegal considerations
48012	for physicians treating patients with depression in the age of SSRIs.
	eMedicine Feature Series, Series 2: Issue 12.
77124	Prescrire International (2014). Mefloquine: persistent vestibular disorders.
11124	Prescrire International, 23(150): 157.
53734	Preti A (2002). Seasonal variation and meteotropism in suicide: clinical
	relevance of findings and implications for research. Acta
	Neuropsychiatrica, 14: 17-28.
53736	Preti A, Miotto P, De Coppi M (2000). Season and suicide: recent
	findings from Italy. Crisis, 21(2): 59-70.
54281	Pridmore S, McArthur M (2009). Suicide and Western culture. Australas
34201	Psychiatry, 17(1): 42-50.
	Prosser JM, Perrone J, Pines JM (2007). The epidemiology of intentional
52681	non-fatal self-harm poisoning in the United States: 2001-2004. J Med
	Toxicol, 3(1): 20-4.
	Quine S, Bernard D, Booth M, et al (2003). Health and access issues
53443	among Australian adolescents: a rural-urban comparison. Retrieved 24
33443	July 2009, from
	http://www.rrh.org.au/publishedarticles/article_print_245.pdf
	Quinn JC (2015). Complex membrane channel blockade: a unifying
76939	hypothesis for the prodromal and acute neuropsychiatric sequelae
70939	resulting from exposure to the antimalarial drug mefloquine. Journal of
	Parasitology Research, ID 368064: 12 pages.
	Rademaker M (2010). Adverse effects of isotretinoin: a retrospective
77964	review of 1743 patients started on isotretinoin. Australian Journal of
	Dermatology, 51(4): 248-53.
	Raguraman J, Vijaysagar KJ, Chandrasekaran R (2004). [Comment] An
87647	unusual presentation of PTSD. Australian & New Zealand Journal of
	Psychiatry, 38(9): 760.
	Rajapakse S, Rodrigo C, Fernando SD (2015). Tafenoquine for
77934	preventing relapse in people with plasmodium vivax malaria (review). The
	Cochrane Collaboration.
	Rajkumar RP (2015). Correlates of suicide-related ideations and attempts
77997	in patients with acute and transient psychotic disorder. Clin Schizophr
	Relat Psychoses, [Epub ahead of print]. [Abstract]
	Ramrakha S, Fergusson DM, Horwood LJ, et al (2015). Cumulative
78063	mental health consequences of acne: 23-year follow-up in general
	population birth cohort study. Br J Dermatol, [Epub ahead of print].
	Ratcliffe GE, Enns MW, Belik SL, (2008). Chronic pain conditions and
54008	suicidal ideation and suicide attempts: an epidemiologic perspective. Clin
	J Pain, 24(3): 204-10.
E4000	Reeves RR, Ladner ME (2009). Antidepressant-induced suicidality:
54023	implications for clinical practice. South Med J, 102(7): 713-8.
	Regambal MJ, Alden LE, Wagner SL et al (2015). Characteristics of the
85927	traumatic stressors experienced by rural first responders. Journal of
33321	Anxiety Disorders, 34: 86-93.
	Reger MA, Smolenski DJ, Skopp NA, et al (2015). Risk of suicide among
79841	US military service members following Operation Enduring Freedom or
	Operation Iraqi Freedom deployment and separation from the US
	military. JAMA Psychiatry, 72(6): 561-9.
	Reich J (1998). The relationship of suicide attempts, borderline
53447	personality traits, and major depressive disorder in a veteran outpatient
	population. J Affect Disord, 49: 151-6.
F.1000	Reith DM, Edmonds L (2007). Assessing the role of drugs in suicidal
54033	ideation and suicidality. CNS Drugs, 21(6): 463-72.
l	

Remafedi G (1999). [Comment] Suicide and sexual orientation. Nearing the end of controversy? Arch Gen Psychiatry, 56: 885-6. Comment on ID: 53682.
Rezaeian M (2008-2009). Islam and suicide: A short personal communication. OMEGA, 58(1): 77-85.
Ribai P, Nguyen K, Hahn-Barma V, et al (2007). Psychiatric and cognitive difficulties as indicators of juvenile Huntington disease onset in 29 patients. Arch Neurol, 64: 813-9.
Richardson AS, Bergen HA, Martin G, et al (1004). Perceived academic performance as an indicator of risk of attempted suicide in young adolescents. Arch Suicide Res, 9(2): 163-76.
Richardson AS, Bergen HA, Martin G, et al (2005). Perceived academic performance as an indicator or risk of attempted suicide in young adolescents. Arch Suicide Res, 9: 163-76.
Ringqvist A, Bech P, Glenthoj B, et al (2015). Acute and long-term psychiatric side effects of mefloquine: A follow-up on Danish adverse event reports. Travel Medicine and Infectious Disease, 13: 80-8.
Rish BL, Dilustro JF, Salazar AM, et al (1997). Spinal cord injury: A 25 year morbidity and mortality study. Military Medicine, 162(2): 141-8.
Ritchie EC, Block J, Nevin RL (2013). Psychiatric side effects of Mefloquine: applications to forensic psychiatry. J Am Acad Psychiatry Law, 41: 224-35.
Robertson HT, Allison DB (2009). Drugs associated with more suicidal ideations are also associated with more suicide attempts. PLoS One, 4(10): e7312.
Roche (2014). Larium. Roche Products.
Rock D, Greenberg D, Hallmayer J (2006). Season-of-birth as a risk factor for the seasonality of suicidal behaviour. Eur Arch Psychiatry Clin Neurosci, 256: 98-105.
Rock DJ, Hallmayer JF (2008). The seasonal risk for deliberate self-harm. Determined by place of birth, but occurrence determined by place of residence. Crisis, 29(4): 191-201.
Rodriguez-Cano T, Beato-Fernandez L, Llario AB (2006). Body dissatisfaction as a predictor of self-reported suicide attempts in adolescents: A Spanish community prospective study. J Adolesc Health, 38: 684-8.
Rogers ML, Kelliher-Rabon J, Hagan CR, et al (2017). Negative emotions in veterans relate to suicide risk through feelings of perceived burdensomeness and thwarted belongingness. J Affect Disord, [Epub ahead of volume]:
Ross EA, Reisfield GM, Watson MC, et al (2012). Psychoactive "bath salts" intoxication with methlenedioxypyrovalerone. Am J Med, 125(9): 854-8.
Rossow I, Lauritzen G (2001). Shattered childhood: a key issue in suicidal behavior among drug addicts? Addiction, 96: 227-40.
Rostila M, Saarela J, Kawachi I (2012). Mortality in parents following the death of a child: a nationwide follow-up study from Sweden. J Epidemiol Comm Health, 66: 927-33.
Rostila M, Saarela J, Kawachi I (2013). Suicide following the death of a sibling: a nationwide follow-up study from Sweden. BMJ Open, 3: e002618.
Rowe JL, Bruce ML, Conwell Y (2006). Correlates of suicide among home health care utilizers who dies by suicide and community controls. Suicide & Life-Threatening Behaviour, 36(1): 65-75.
Roy A (1993). Anxiety and suicide in depression. Can J Psychiatry, 38(10): 694-5.

Roy A, Karoum F, Pollack S (1992). Marked reduction in indexes of dopamine metabolism among patients with depression who attempt suicide. Arch Gen Psychiatry, 49: 447-50.  Rozanov V, Caril V (2012). Suicide among war veterans. Int J Environ Res Public Health, 9: 2504-19.  Rubenowitz E, Waern M, Wilhelmson K, et al (2001). Life events and psychosocial factors in elderly suicides - a case-control study. Psychol Med, 31: 1193-202.  Rudatsikira E, Muula AS, Siziya S, et al (2007). Suicidal ideation and associated factors among school-going adolescents in rural Uganda. BMC Psychiatry, 7: 67.  Rudd MD, Dahm PF, Rajab MH (1993). Diagnostic comorbidity in persons with suicidal ideation and behaviour. Am J Psychiatry, 150(6): 928-34.  1770 Runeson BS (1989). Mental disorder in youth suicide: DSM-III-R Axes I and II. Acta Psychiatr Scand, 79: 490-7.  Ryb GE, Soderstrom CA, Kufera JA, et al (2006). Longitudinal study of suicide after traumatic injury. J Trauma, 61: 799-804.  Ryding E, Ahnilde JA, Lindstrom M, et al (2006). Regional brain serotonin and dopamine transporter binding capacity in suicide attempters relate to impulsiveness and mental energy. Psychiatry Res Neuroimaging, 148: 195-203.  Ryding E, Lindstrom M, Traskman-Bendz L (2008). The role of dopamine and serotonin in suicidal behaviour and aggression. Progress in Brain Research, Chapter 15, 172: 307-15.  Sahin HA, Sahin I, Arabaci F (2003). Sociodemographic factors in organophosphate poisonings: a prospective study. Human & Experimental Toxicology, 22: 349-53.  Sakuta T (1995). A study of murder followed by suicide. Med Law, 14: 141-53.  Sakuta T (1995). A study of murder followed by suicide. Med Law, 14: 141-53.  Sareen J, Cox BJ, Affif TO, et al (2005). Anxiety disorders and risk for suicideal med for mental health care. Arch Gen Psychiatry, 64(7): 843-52.  Schaffer LC, Schaffer CB, Hunter S, et al (2010). Psychiatric reactions to isotretinoin in patients with bipolar disorder. Journal of Affective Disorder, 122(3): 306-8.  Scheider B, Wetterling T, Sargk D	F	
79681 Rozanov V, Carli V (2012). Suicide among war veterans. Int J Environ Res Public Health, 9: 2504-19. Rubenowitz E, Waern M, Wilhelmson K, et al (2001). Life events and psychosocial factors in elderly suicides - a case-control study. Psychol Med, 31: 1193-202. Rudatsikira E, Muula AS, Siziya S, et al (2007). Suicidal ideation and associated factors among school-going adolescents in rural Uganda. BMC Psychiatry, 7: 67. Rudd MD, Dahm PF, Rajab MH (1993). Diagnostic comorbidity in persons with suicidal ideation and behaviour. Am J Psychiatry, 150(6): 928-34.  Runeson BS (1989). Mental disorder in youth suicide: DSM-III-R Axes I and II. Acta Psychiatr Scand, 79: 490-7.  Ryb GE, Soderstrom CA, Kufera JA, et al (2006). Longitudinal study of suicide after traumatic injury. J Trauma, 61: 799-804.  Ryding E, Ahnilde JA, Lindstrom M, et al (2006). Regional brain serotonin and dopamine transporter binding capacity in suicide attempters relate to impulsiveness and mental energy. Psychiatry Res Neuroimaging, 148: 195-203.  Ryding E, Lindstrom M, Traskman-Bendz L (2008). The role of dopamine and serotonin in suicidal behaviour and aggression. Progress in Brain Research, Chapter 15, 172: 307-15.  Sahin HA, Sahin I, Arabaci F (2003). Sociodemographic factors in organophosphate poisonings: a prospective study. Human & Experimental Toxicology, 22: 349-53.  Sakinofskyl (2003). Suicide: the persisting challenge. Can J Psychiatry, 48(5): 289-91.  Sakinofskyl (2003). Suicide: the persisting challenge. Can J Psychiatry, 48(5): 289-91.  Santiago JM, Dalen JE (1994). Cholesterol and violent behaviour. Arch Intern Med, 154: 1317-21.  Sareen J, Belik S-L (2009). [Comment] The need for outreach in preventing suicide among young veterans. PLoS Med, 6(3): e1000035. Comment on ID: 53445.  Sareen J, Cox BJ, Affif TO, et al (2007). Anxiety disorders and risk for suicidal ideation and suicide attempts. Arch Gen Psychiatry, 62: 1249-57.  Sareen J, Cox BJ, Affif TO, et al (2007). Combat and peacekeeping operations in relation to prevalence of	1769	Roy A, Karoum F, Pollack S (1992). Marked reduction in indexes of dopamine metabolism among patients with depression who attempt
Res Public Health, 9: 2504-19.  Rubenowitz E, Waern M, Wilhelmson K, et al (2001). Life events and psychosocial factors in elderly suicides - a case-control study. Psychol Med, 31: 1193-202.  Rudatsikria E, Muula AS, Siziya S, et al (2007). Suicidal ideation and associated factors among school-going adolescents in rural Uganda. BMC Psychiatry, 7: 67.  Rudd MD, Dahm PF, Rajab MH (1993). Diagnostic comorbidity in persons with suicidal ideation and behaviour. Am J Psychiatry, 150(6): 928-34.  Runeson BS (1989). Mental disorder in youth suicide: DSM-III-R Axes I and II. Acta Psychiatr Scand, 79: 490-7.  Ryb GE, Soderstrom CA, Kufera JA, et al (2006). Longitudinal study of suicide after traumatic injury. J Trauma, 61: 799-804.  Ryding E, Ahnilde JA, Lindstrom M, et al (2006). Regional brain serotonin and dopamine transporter binding capacity in suicide atmepters relate to impulsiveness and mental energy. Psychiatry Res Neuroimaging, 148: 195-203.  Ryding E, Lindstrom M, Traskman-Bendz L (2008). The role of dopamine and serotonin in suicidal behaviour and aggression. Progress in Brain Research, Chapter 15, 172: 307-15.  Sahin HA, Sahin I, Arabaci F (2003). Sociodemographic factors in organophosphate poisonings: a prospective study. Human & Experimental Toxicology, 22: 349-53.  Sakinofsky I (2003). Suicide: the persisting challenge. Can J Psychiatry, 48(5): 289-91.  Sakuta T (1995). A study of murder followed by suicide. Med Law, 14: 141-53.  Sanofi-Aventis Australia (2012). Plaquenil. Product Information.  Santiago JM, Dalen JE (1994). Cholesterol and violent behaviour. Arch Intern Med, 154: 1317-21.  Sareen J, Belik S-L (2009). [Comment] The need for outreach in preventing suicide among young veterans. PLoS Med, 6(3): e1000035. Comment on ID: 53445.  Sareen J, Cox BJ, Affif TO, et al (2005). Anxiety disorders and risk for suicidal ideation and suicide attempts. Arch Gen Psychiatry, 62: 1249-57.  Sareen J, Cox BJ, Affif TO, et al (2007). Combat and peacekeeping operations in relation to prevalence of mental disorders		
Rubenowitz E, Waern M, Wilhelmson K, et al (2001). Life events and psychosocial factors in elderly suicides - a case-control study. Psychol Med, 31: 1193-202. Rudatsikira E, Muula AS, Siziya S, et al (2007). Suicidal ideation and associated factors among school-going adolescents in rural Uganda. BMC Psychiatry, 7: 67. Rudd MD, Dahm PF, Rajab MH (1993). Diagnostic comorbidity in persons with suicidal ideation and behaviour. Am J Psychiatry, 150(6): 928-34.  Runeson BS (1989). Mental disorder in youth suicide: DSM-III-R Axes I and II. Acta Psychiatr Scand, 79: 490-7.  Ryb GE, Soderstrom CA, Kufera JA, et al (2006). Longitudinal study of suicide after traumatic injury. J Trauma, 61: 799-804.  Ryding E, Ahnlide JA, Lindstrom M, et al (2006). Regional brain serotonin and dopamine transporter binding capacity in suicide attempters relate to impulsiveness and mental energy. Psychiatry Res Neuroimaging, 148: 195-203.  Ryding E, Lindstrom M, Traskman-Bendz L (2008). The role of dopamine and serotonin in suicidal behaviour and aggression. Progress in Brain Research, Chapter 15, 172: 307-15.  Sahin HA, Sahin I, Arabaci F (2003). Sociodemographic factors in organophosphate poisonings: a prospective study. Human & Experimental Toxicology, 22: 349-53.  Sakinofsky I (2003). Suicide: the persisting challenge. Can J Psychiatry, 48(5): 289-91.  Sakuta T (1995). A study of murder followed by suicide. Med Law, 14: 141-53.  Sancii-Aventis Australia (2012). Plaquenil. Product Information.  Santiago JM, Dalen JE (1994). Cholesterol and violent behaviour. Arch Intern Med, 154: 1317-21.  Sareen J, Cox BJ, Affif TO, et al (2005). Anxiety disorders and risk for suicidal ideation and suicide attempts. Arch Gen Psychiatry, 62: 1249-57.  Sareen J, Cox BJ, Affif TO, et al (2007). Combat and peacekeeping operations in relation to prevalence of mental disorders and perceived need for mental health care. Arch Gen Psychiatry, 62: 1249-57.  Schoenbaum M, Kessler RC, Gilman SE, et al (2014). Predictors of suicide and accident death in the Army Stu	79681	
54766 psychosocial factors in elderly suicides - a case-control study. Psychol Med, 31: 1193-202.  Rudatsikira E, Muula AS, Siziya S, et al (2007). Suicidal ideation and associated factors among school-going adolescents in rural Uganda. BMC Psychiatry, 7: 67.  Rudd MD, Dahm PF, Rajab MH (1993). Diagnostic comorbidity in persons with suicidal ideation and behaviour. Am J Psychiatry, 150(6): 928-34.  1770 Rueson BS (1989). Mental disorder in youth suicide: DSM-III-R Axes I and II. Acta Psychiatr Scand, 79: 490-7.  Ryb GE, Soderstrom CA, Kufera JA, et al (2006). Longitudinal study of suicide affer traumatic injury. J Trauma, 61: 799-804.  Ryding E, Ahnlide JA, Lindstrom M, et al (2006). Regional brain serotonin and dopamine transporter binding capacity in suicide attempters relate to impulsiveness and mental energy. Psychiatry Res Neuroimaging, 148: 195-203.  Ryding E, Lindstrom M, Traskman-Bendz L (2008). The role of dopamine and serotonin in suicidal behaviour and aggression. Progress in Brain Research, Chapter 15, 172: 307-15.  Sahin HA, Sahin I, Arabaci F (2003). Sociodemographic factors in organophosphate poisonings: a prospective study. Human & Experimental Toxicology, 22: 349-53.  Sakinofisky I (2003). Suicide: the persisting challenge. Can J Psychiatry, 48(5): 289-91.  Sakina T (1995). A study of murder followed by suicide. Med Law, 14: 141-53.  78683 Sanofi-Aventis Australia (2012). Plaquenil. Product Information.  Sareen J, Belik S-L (2009). [Comment] The need for outreach in preventing suicide among young veterans. PLoS Med, 6(3): e1000035. Comment on ID: 53445.  Sareen J, Cox BJ, Affif TO, et al (2005). Anxiety disorders and risk for suicidal ideation and suicide attempts. Arch Gen Psychiatry, 64(7): 843-52.  Schaffer LC, Schaffer CB, Hunter S, et al (2010). Psychiatry 64(7): 843-52.  Schenèder B, Wetterling T, Sargk D, et al (2006). Axis I disorders and personality disorders as risk factors for suicide. Eur Arch Psychiatry Clin Neurosci, 256: 17-27.  Schoenbaum M, Kessler RC, Gilman SE, et al (2014).	54766	
S2468 Rudatsikira E, Muula AS, Siziya S, et al (2007). Suicidal ideation and associated factors among school-going adolescents in rural Uganda. BMC Psychiatry, 7: 67.  Rudd MD, Dahm PF, Rajab MH (1993). Diagnostic comorbidity in persons with suicidal ideation and behaviour. Am J Psychiatry, 150(6): 928-34.  1770 Runeson BS (1989). Mental disorder in youth suicide: DSM-III-R Axes I and II. Acta Psychiatr Scand, 79: 490-7.  Ryb GE, Soderstrom CA, Kufera JA, et al (2006). Longitudinal study of suicide after traumatic injury. J Trauma, 61: 799-804.  Ryding E, Ahnilde JA, Lindstrom M, et al (2006). Regional brain serotonin and dopamine transporter binding capacity in suicide attempters relate to impulsiveness and mental energy. Psychiatry Res Neuroimaging, 148: 195-203.  Ryding E, Lindstrom M, Traskman-Bendz L (2008). The role of dopamine and serotonin in suicidal behaviour and aggression. Progress in Brain Research, Chapter 15, 172: 307-15.  Sahin HA, Sahin I, Arabaci F (2003). Sociodemographic factors in organophosphate poisonings: a prospective study. Human & Experimental Toxicology, 22: 349-53.  Sakinofsky I (2003). Suicide: the persisting challenge. Can J Psychiatry, 48(5): 289-91.  Sakinofsky I (2003). Suicide: the persisting challenge. Can J Psychiatry, 48(5): 289-91.  Sakinofsky I (2003). Suicide: the persisting challenge. Can J Psychiatry, 48(5): 289-91.  Sariago JM, Dalen JE (1994). Cholesterol and violent behaviour. Arch Intern Med, 154: 1317-21.  Sareen J, Bart S, L (2009). [Comment] The need for outreach in preventing suicide among young veterans. PLoS Med, 6(3): e1000035. Comment on ID: 53445.  Sareen J, Cox BJ, Affif TO, et al (2005). Anxiety disorders and risk for suicidal ideation and suicide attempts. Arch Gen Psychiatry, 62: 1249-57.  Sareen J, Cox BJ, Affif TO, et al (2005). Combat and peacekeeping operations in relation to prevalence of mental disorders and perceived need for mental health care. Arch Gen Psychiatry, 64(7): 843-52.  Schaffer LC, Schaffer CB, Hunter S, et al (2010). Psychiatric re		
ssociated factors among school-going adolescents in rural Uganda. BMC Psychiatry, 7: 67.  Rudd MD, Dahm PF, Rajab MH (1993). Diagnostic comorbidity in persons with suicidal ideation and behaviour. Am J Psychiatry, 150(6): 928-34.  Runeson BS (1989). Mental disorder in youth suicide: DSM-III-R Axes I and II. Acta Psychiatr Scand, 79: 490-7.  Ryb GE, Soderstrom CA, Kufera JA, et al (2006). Longitudinal study of suicide after traumatic injury. J Trauma, 61: 799-804.  Ryding E, Ahnlide JA, Lindstrom M, et al (2006). Regional brain serotonin and dopamine transporter binding capacity in suicide attempters relate to impulsiveness and mental energy. Psychiatry Res Neuroimaging, 148: 195-203.  Ryding E, Lindstrom M, Traskman-Bendz L (2008). The role of dopamine and serotonin in suicidal behaviour and aggression. Progress in Brain Research, Chapter 15, 172: 307-15.  Sahin HA, Sahin I, Arabaci F (2003). Sociodemographic factors in organophosphate poisonings: a prospective study. Human & Experimental Toxicology, 22: 349-53.  Sakuta T (1995). A study of murder followed by suicide. Med Law, 14: 141-53.  Sakuta T (1995). A study of murder followed by suicide. Med Law, 14: 141-53.  Sanofi-Aventis Australia (2012). Plaquenil. Product Information.  Santiago JM, Dalen JE (1994). Cholesterol and violent behaviour. Arch Intern Med., 154: 1317-21.  Sareen J, Belik S-L (2009). [Comment] The need for outreach in preventing suicide among young veterans. PLoS Med, 6(3): e1000035. Comment on ID: 53445.  Sareen J, Cox BJ, Affif TO, et al (2007). Combat and peacekeeping operations in relation to prevalence of mental disorders and perceived need for mental health care. Arch Gen Psychiatry, 62: 1249-57.  Schoerbaum M, Kessler RC, Gilman SE, et al (2014). Predictors of suicide and accident death in the Army Study to Assess Risk and Personality disorders as risk factors for suicide. Eur Arch Psychiatry Clin Neurosci, 256: 17-27.  Schoenbaum M, Kessler RC, Gilman SE, et al (2014). Predictors of suicide and accident death in the Army Study to As		Med, 31: 1193-202.
Rudd MD, Dahm PF, Rajab MH (1993). Diagnostic comorbidity in persons with suicidal ideation and behaviour. Am J Psychiatry, 150(6): 928-34.  1770 Runeson BS (1989). Mental disorder in youth suicide: DSM-III-R Axes I and II. Acta Psychiatr Scand, 79: 490-7.  Ryb GE, Soderstrom CA, Kufera JA, et al (2006). Longitudinal study of suicide after traumatic injury. J Trauma, 61: 799-804.  Ryding E, Ahnlide JA, Lindstrom M, et al (2006). Regional brain serotonin and dopamine transporter binding capacity in suicide attempters relate to impulsiveness and mental energy. Psychiatry Res Neuroimaging, 148: 195-203.  Ryding E, Lindstrom M, Traskman-Bendz L (2008). The role of dopamine and serotonin in suicidal behaviour and aggression. Progress in Brain Research, Chapter 15, 172: 307-15.  Sahin HA, Sahin I, Arabaci F (2003). Sociodemographic factors in organophosphate poisonings: a prospective study. Human & Experimental Toxicology, 22: 349-53.  Sakinofsky I (2003). Suicide: the persisting challenge. Can J Psychiatry, 48(5): 289-91.  Sakuta T (1995). A study of murder followed by suicide. Med Law, 14: 141-53.  78683 Sanofi-Aventis Australia (2012). Plaquenil. Product Information.  4194 Santiago JM, Dalen JE (1994). Cholesterol and violent behaviour. Arch Intern Med, 154: 1317-21.  Sareen J, Belik S-L (2009). [Comment] The need for outreach in preventing suicide among young veterans. PLoS Med, 6(3): e1000035. Comment on ID: 53445.  Sareen J, Cox BJ, Affif TO, et al (2005). Anxiety disorders and risk for suicidal ideation and suicide attempts. Arch Gen Psychiatry, 62: 1249-57.  Sareen J, Cox BJ, Affif TO, et al (2007). Combat and peacekeeping operations in relation to prevalence of mental disorders and perceived need for mental health care. Arch Gen Psychiatry, 62: 1249-57.  Sareen J, Cox BJ, Affif TO, et al (2007). Combat and peacekeeping operations in relation to prevalence of mental disorders and perceived need for mental health care. Arch Gen Psychiatry, 64(7): 843-52.  Schaffer LC, Schaffer CB, Hunter S, et al (2014). Psych		
4201 persons with suicidal ideation and behaviour. Am J Psychiatry, 150(6): 928-34.  1770 Runeson BS (1989). Mental disorder in youth suicide: DSM-III-R Axes I and II. Acta Psychiatr Scand, 79: 490-7.  49922 Ryb GE, Soderstrom CA, Kufera JA, et al (2006). Longitudinal study of suicide after traumatic injury. J Trauma, 61: 799-804.  Ryding E, Ahnlide JA, Lindstrom M, et al (2006). Regional brain serotonin and dopamine transporter binding capacity in suicide attempters relate to impulsiveness and mental energy. Psychiatry Res Neuroimaging, 148: 195-203.  Ryding E, Lindstrom M, Traskman-Bendz L (2008). The role of dopamine and serotonin in suicidal behaviour and aggression. Progress in Brain Research, Chapter 15, 172: 307-15.  Sahin HA, Sahin I, Arabaci F (2003). Sociodemographic factors in organophosphate poisonings: a prospective study. Human & Experimental Toxicology, 22: 349-53.  Sakinofsky I (2003). Suicide: the persisting challenge. Can J Psychiatry, 48(5): 289-91.  Sakuta T (1995). A study of murder followed by suicide. Med Law, 14: 141-53.  78683 Sanofi-Aventis Australia (2012). Plaquenil. Product Information.  Santiago JM, Dalen JE (1994). Cholesterol and violent behaviour. Arch Intern Med, 154: 1317-21.  Sareen J, Belik S-L (2009). [Comment] The need for outreach in preventing suicide among young veterans. PLoS Med, 6(3): e1000035. Comment on ID: 53445.  Sareen J, Cox BJ, Affif TO, et al (2005). Anxiety disorders and priceived need for mental health care. Arch Gen Psychiatry, 62: 1249-57.  Sareen J, Cox BJ, Affif TO, et al (2007). Combat and peacekeeping operations in relation to prevalence of mental disorders and perceived need for mental health care. Arch Gen Psychiatry, 64(7): 843-52.  Schaffer LC, Schaffer CB, Hunter S, et al (2010). Psychiatric reactions to isotretinoin in patients with bipolar disorder. Journal of Affective Disorder, 122(3): 306-8.  Schneider B, Wetterling T, Sargk D, et al (2006). Axis I disorders and perceived need for mental disorders as risk factors for suicide. Eur Arch Psychi	52468	
928-34.  1770 Runeson BS (1989). Mental disorder in youth suicide: DSM-III-R Axes I and II. Acta Psychiatr Scand, 79: 490-7.  Ryb GE, Soderstrom CA, Kufera JA, et al (2006). Longitudinal study of suicide after traumatic injury. J Trauma, 61: 799-804.  Ryding E, Ahnlide JA, Lindstrom M, et al (2006). Regional brain serotonin and dopamine transporter binding capacity in suicide attempters relate to impulsiveness and mental energy. Psychiatry Res Neuroimaging, 148: 195-203.  Ryding E, Lindstrom M, Traskman-Bendz L (2008). The role of dopamine and serotonin in suicidal behaviour and aggression. Progress in Brain Research, Chapter 15, 172: 307-15.  Sahin HA, Sahin I, Arabaci F (2003). Sociodemographic factors in organophosphate poisonings: a prospective study. Human & Experimental Toxicology, 22: 349-53.  Sakinofsky I (2003). Suicide: the persisting challenge. Can J Psychiatry, 48(5): 289-91.  52439 Sakuta T (1995). A study of murder followed by suicide. Med Law, 14: 141-53.  78683 Sanofi-Aventis Australia (2012). Plaquenil. Product Information.  Santiago JM, Dalen JE (1994). Cholesterol and violent behaviour. Arch Intern Med, 154: 1317-21.  Sareen J, Belik S-L (2009). [Comment] The need for outreach in preventing suicide among young veterans. PLoS Med, 6(3): e1000035. Comment on ID: 53445.  Sareen J, Cox BJ, Affif TO, et al (2005). Anxiety disorders and risk for suicidal ideation and suicide attempts. Arch Gen Psychiatry, 62: 1249-57.  Sareen J, Cox BJ, Affif TO, et al (2007). Combat and peacekeeping operations in relation to prevalence of mental disorders and personality disorders as risk factors for suicide. Eur Arch Psychiatry Clin Neurosci, 256: 47-27.  Schoenbaum M, Kessler RC, Gilman SE, et al (2010). Psychiatric reactions to siotretinoin in patients with bipolar disorder. Journal of Affective Disorder, 122(3): 306-8.  Schneider B, Wetterling T, Sargk D, et al (2006). Axis I disorders and personality disorders as risk factors for suicide. Eur Arch Psychiatry Clin Neurosci, 256: 47-27.  Schoenbaum M, Kessler R		
and II. Acta Psychiatr Scand, 79: 490-7.  Ryb GE, Soderstrom CA, Kufera JA, et al (2006). Longitudinal study of suicide after traumatic injury. J Trauma, 61: 799-804.  Ryding E, Ahnlide JA, Lindstrom M, et al (2006). Regional brain serotonin and dopamine transporter binding capacity in suicide attempters relate to impulsiveness and mental energy. Psychiatry Res Neuroimaging, 148: 195-203.  Ryding E, Lindstrom M, Traskman-Bendz L (2008). The role of dopamine and serotonin in suicidal behaviour and aggression. Progress in Brain Research, Chapter 15, 172: 307-15.  Sahin HA, Sahin I, Arabaci F (2003). Sociodemographic factors in organophosphate poisonings: a prospective study. Human & Experimental Toxicology, 22: 349-53.  Sakinofsky I (2003). Suicide: the persisting challenge. Can J Psychiatry, 48(5): 289-91.  Sakuta T (1995). A study of murder followed by suicide. Med Law, 14: 141-53.  78683 Sanofi-Aventis Australia (2012). Plaquenil. Product Information.  Santiago JM, Dalen JE (1994). Cholesterol and violent behaviour. Arch Intern Med, 154: 1317-21.  Sareen J, Belik S-L (2009). [Comment] The need for outreach in preventing suicide among young veterans. PLoS Med, 6(3): e1000035. Comment on ID: 53445.  Sareen J, Cox BJ, Afffi TO, et al (2005). Anxiety disorders and risk for suicidal ideation and suicide attempts. Arch Gen Psychiatry, 62: 1249-57.  Sareen J, Cox BJ, Afffi TO, et al (2007). Combat and peacekeeping operations in relation to prevalence of mental disorders and perceived need for mental health care. Arch Gen Psychiatry, 64(7): 843-52.  Schaffer LC, Schaffer CB, Hunter S, et al (2010). Psychiatric reactions to isotretinoin in patients with bipolar disorder. Journal of Affective Disorder, 122(3): 306-8.  Schneider B, Wetterling T, Sargk D, et al (2006). Axis I disorders and personality disorders as risk factors for suicide. Eur Arch Psychiatry Clin Neurosci, 256: 17-27.  Schoenbaum M, Kessler RC, Gilman SE, et al (2014). Predictors of suicide and accident death in the Army Study to Assess Risk and Resilien	4201	
and II. Acta Psychiatr Scand, 79: 490-7.  Ryb GE, Soderstrom CA, Kufera JA, et al (2006). Longitudinal study of suicide after traumatic injury. J Trauma, 61: 799-804.  Ryding E, Ahnlide JA, Lindstrom M, et al (2006). Regional brain serotonin and dopamine transporter binding capacity in suicide attempters relate to impulsiveness and mental energy. Psychiatry Res Neuroimaging, 148: 195-203.  Ryding E, Lindstrom M, Traskman-Bendz L (2008). The role of dopamine and serotonin in suicidal behaviour and aggression. Progress in Brain Research, Chapter 15, 172: 307-15.  Sahin HA, Sahin I, Arabaci F (2003). Sociodemographic factors in organophosphate poisonings: a prospective study. Human & Experimental Toxicology, 22: 349-53.  Sakinofsky I (2003). Suicide: the persisting challenge. Can J Psychiatry, 48(5): 289-91.  Sakuta T (1995). A study of murder followed by suicide. Med Law, 14: 141-53.  78683 Sanofi-Aventis Australia (2012). Plaquenil. Product Information.  Santiago JM, Dalen JE (1994). Cholesterol and violent behaviour. Arch Intern Med, 154: 1317-21.  Sareen J, Belik S-L (2009). [Comment] The need for outreach in preventing suicide among young veterans. PLoS Med, 6(3): e1000035. Comment on ID: 53445.  Sareen J, Cox BJ, Afffi TO, et al (2005). Anxiety disorders and risk for suicidal ideation and suicide attempts. Arch Gen Psychiatry, 62: 1249-57.  Sareen J, Cox BJ, Afffi TO, et al (2007). Combat and peacekeeping operations in relation to prevalence of mental disorders and perceived need for mental health care. Arch Gen Psychiatry, 64(7): 843-52.  Schaffer LC, Schaffer CB, Hunter S, et al (2010). Psychiatric reactions to isotretinoin in patients with bipolar disorder. Journal of Affective Disorder, 122(3): 306-8.  Schneider B, Wetterling T, Sargk D, et al (2006). Axis I disorders and personality disorders as risk factors for suicide. Eur Arch Psychiatry Clin Neurosci, 256: 17-27.  Schoenbaum M, Kessler RC, Gilman SE, et al (2014). Predictors of suicide and accident death in the Army Study to Assess Risk and Resilien	4770	
suicide after traumatic injury. J Trauma, 61: 799-804.  Ryding E, Ahnlide JA, Lindstrom M, et al (2006). Regional brain serotonin and dopamine transporter binding capacity in suicide attempters relate to impulsiveness and mental energy. Psychiatry Res Neuroimaging, 148: 195-203.  Ryding E, Lindstrom M, Traskman-Bendz L (2008). The role of dopamine and serotonin in suicidal behaviour and aggression. Progress in Brain Research, Chapter 15, 172: 307-15.  Sahin HA, Sahin I, Arabaci F (2003). Sociodemographic factors in organophosphate poisonings: a prospective study. Human & Experimental Toxicology, 22: 349-53.  Sakinofsky I (2003). Suicide: the persisting challenge. Can J Psychiatry, 48(5): 289-91.  Sakuta T (1995). A study of murder followed by suicide. Med Law, 14: 141-53.  Sanofi-Aventis Australia (2012). Plaquenil. Product Information.  Santiago JM, Dalen JE (1994). Cholesterol and violent behaviour. Arch Intern Med, 154: 1317-21.  Sareen J, Belik S-L (2009). [Comment] The need for outreach in preventing suicide among young veterans. PLoS Med, 6(3): e1000035. Comment on ID: 53445.  Sareen J, Cox BJ, Affif TO, et al (2005). Anxiety disorders and risk for suicidal ideation and suicide attempts. Arch Gen Psychiatry, 62: 1249-57.  Sareen J, Cox BJ, Affif TO, et al (2007). Combat and peacekeeping operations in relation to prevalence of mental disorders and perceived need for mental health care. Arch Gen Psychiatry, 64(7): 843-52.  Schaffer LC, Schaffer CB, Hunter S, et al (2010). Psychiatric reactions to isotretinoin in patients with bipolar disorder. Journal of Affective Disorder, 122(3): 306-8.  Schneider B, Wetterling T, Sargk D, et al (2006). Axis I disorders and personality disorders as risk factors for suicide. Eur Arch Psychiatry Clin Neurosci, 256: 17-27.  Schoenbaum M, Kessler RC, Gilman SE, et al (2014). Predictors of suicide and accident death in the Army Study to Assess Risk and Resilience in Servicemembers (Army	1770	and II. Acta Psychiatr Scand, 79: 490-7.
Ryding E, Ahnlide JA, Lindstrom M, et al (2006). Regional brain serotonin and dopamine transporter binding capacity in suicide attempters relate to impulsiveness and mental energy. Psychiatry Res Neuroimaging, 148: 195-203.  Ryding E, Lindstrom M, Traskman-Bendz L (2008). The role of dopamine and serotonin in suicidal behaviour and aggression. Progress in Brain Research, Chapter 15, 172: 307-15.  Sahin HA, Sahin I, Arabaci F (2003). Sociodemographic factors in organophosphate poisonings: a prospective study. Human & Experimental Toxicology, 22: 349-53.  54029 Sakinofsky I (2003). Suicide: the persisting challenge. Can J Psychiatry, 48(5): 289-91.  Sakuta T (1995). A study of murder followed by suicide. Med Law, 14: 141-53.  78683 Sanofi-Aventis Australia (2012). Plaquenil. Product Information.  Santiago JM, Dalen JE (1994). Cholesterol and violent behaviour. Arch Intern Med, 154: 1317-21.  Sareen J, Belik S-L (2009). [Comment] The need for outreach in preventing suicide among young veterans. PLoS Med, 6(3): e1000035. Comment on ID: 53445.  Sareen J, Cox BJ, Affif TO, et al (2005). Anxiety disorders and risk for suicidal ideation and suicide attempts. Arch Gen Psychiatry, 62: 1249-57.  Sareen J, Cox BJ, Affif TO, et al (2007). Combat and peacekeeping operations in relation to prevalence of mental disorders and perceived need for mental health care. Arch Gen Psychiatry, 64(7): 843-52.  Schaffer LC, Schaffer CB, Hunter S, et al (2010). Psychiatric reactions to isotretinoin in patients with bipolar disorder. Journal of Affective Disorder, 122(3): 306-8.  Schneider B, Wetterling T, Sargk D, et al (2006). Axis I disorders and personality disorders as risk factors for suicide. Eur Arch Psychiatry Clin Neurosci, 256: 17-27.  Schoenbaum M, Kessler RC, Gilman SE, et al (2014). Predictors of suicide and accident death in the Army Study to Assess Risk and Resilience in Servicemembers (Army	40022	Ryb GE, Soderstrom CA, Kufera JA, et al (2006). Longitudinal study of
and dopamine transporter binding capacity in suicide attempters relate to impulsiveness and mental energy. Psychiatry Res Neuroimaging, 148: 195-203.  Ryding E, Lindstrom M, Traskman-Bendz L (2008). The role of dopamine and serotonin in suicidal behaviour and aggression. Progress in Brain Research, Chapter 15, 172: 307-15.  Sahin HA, Sahin I, Arabaci F (2003). Sociodemographic factors in organophosphate poisonings: a prospective study. Human & Experimental Toxicology, 22: 349-53.  Sakinofskyl (2003). Suicide: the persisting challenge. Can J Psychiatry, 48(5): 289-91.  Sakuta T (1995). A study of murder followed by suicide. Med Law, 14: 141-53.  Sanofi-Aventis Australia (2012). Plaquenil. Product Information.  Santiago JM, Dalen JE (1994). Cholesterol and violent behaviour. Arch Intern Med, 154: 1317-21.  Sareen J, Belik S-L (2009). [Comment] The need for outreach in preventing suicide among young veterans. PLoS Med, 6(3): e1000035. Comment on ID: 53445.  Sareen J, Cox BJ, Affif TO, et al (2005). Anxiety disorders and risk for suicidal ideation and suicide attempts. Arch Gen Psychiatry, 62: 1249-57.  Sareen J, Cox BJ, Affif TO, et al (2007). Combat and peacekeeping operations in relation to prevalence of mental disorders and perceived need for mental health care. Arch Gen Psychiatry, 64(7): 843-52.  Schaffer LC, Schaffer CB, Hunter S, et al (2010). Psychiatric reactions to isotretinoin in patients with bipolar disorder. Journal of Affective Disorder, 122(3): 306-8.  Schneider B, Wetterling T, Sargk D, et al (2006). Axis I disorders and personality disorders as risk factors for suicide. Eur Arch Psychiatry Clin Neurosci, 256: 17-27.  Schoenbaum M, Kessler RC, Gilman SE, et al (2014). Predictors of suicide and accident death in the Army Study to Assess Risk and Resilience in Servicemembers (Army STARRS): results from the Army Study to Assess Risk and Resilience in Servicemembers (Army	49922	suicide after traumatic injury. J Trauma, 61: 799-804.
impulsiveness and mental energy. Psychiatry Res Neuroimaging, 148: 195-203.  Ryding E, Lindstrom M, Traskman-Bendz L (2008). The role of dopamine and serotonin in suicidal behaviour and aggression. Progress in Brain Research, Chapter 15, 172: 307-15.  Sahin HA, Sahin I, Arabaci F (2003). Sociodemographic factors in organophosphate poisonings: a prospective study. Human & Experimental Toxicology, 22: 349-53.  Sakinofskyl (2003). Suicide: the persisting challenge. Can J Psychiatry, 48(5): 289-91.  Sakuta T (1995). A study of murder followed by suicide. Med Law, 14: 141-53.  Sanofi-Aventis Australia (2012). Plaquenil. Product Information.  Santiago JM, Dalen JE (1994). Cholesterol and violent behaviour. Arch Intern Med, 154: 1317-21.  Sareen J, Belik S-L (2009). [Comment] The need for outreach in preventing suicide among young veterans. PLoS Med, 6(3): e1000035. Comment on ID: 53445.  Sareen J, Cox BJ, Affii TO, et al (2005). Anxiety disorders and risk for suicidal ideation and suicide attempts. Arch Gen Psychiatry, 62: 1249-57.  Sareen J, Cox BJ, Affii TO, et al (2007). Combat and peacekeeping operations in relation to prevalence of mental disorders and perceived need for mental health care. Arch Gen Psychiatry, 64(7): 843-52.  Schaffer LC, Schaffer CB, Hunter S, et al (2010). Psychiatric reactions to isotretinoin in patients with bipolar disorder. Journal of Affective Disorder, 122(3): 306-8.  Schneider B, Wetterling T, Sargk D, et al (2006). Axis I disorders and personality disorders as risk factors for suicide. Eur Arch Psychiatry Clin Neurosci, 256: 17-27.  Schoenbaum M, Kessler RC, Gilman SE, et al (2014). Predictors of suicide and accident death in the Army Study to Assess Risk and Resilience in Servicemembers (Army STARRS): results from the Army Study to Assess Risk and Resilience in Servicemembers (Army		
Impulsiveness and mental energy. Psychiatry Res Neuroimaging, 148: 195-203.  Ryding E, Lindstrom M, Traskman-Bendz L (2008). The role of dopamine and serotonin in suicidal behaviour and aggression. Progress in Brain Research, Chapter 15, 172: 307-15.  Sahin HA, Sahin I, Arabaci F (2003). Sociodemographic factors in organophosphate poisonings: a prospective study. Human & Experimental Toxicology, 22: 349-53.  Sakinofsky I (2003). Suicide: the persisting challenge. Can J Psychiatry, 48(5): 289-91.  Sakuta T (1995). A study of murder followed by suicide. Med Law, 14: 141-53.  78683 Sanofi-Aventis Australia (2012). Plaquenil. Product Information.  Santiago JM, Dalen JE (1994). Cholesterol and violent behaviour. Arch Intern Med, 154: 1317-21.  Sareen J, Belik S-L (2009). [Comment] The need for outreach in preventing suicide among young veterans. PLoS Med, 6(3): e1000035. Comment on ID: 53445.  Sareen J, Cox BJ, Affif TO, et al (2005). Anxiety disorders and risk for suicidal ideation and suicide attempts. Arch Gen Psychiatry, 62: 1249-57.  Sareen J, Cox BJ, Affif TO, et al (2007). Combat and peacekeeping operations in relation to prevalence of mental disorders and perceived need for mental health care. Arch Gen Psychiatry, 64(7): 843-52.  Schaffer LC, Schaffer CB, Hunter S, et al (2010). Psychiatric reactions to isotretinoin in patients with bipolar disorder. Journal of Affective Disorder, 122(3): 306-8.  Schneider B, Wetterling T, Sargk D, et al (2006). Axis I disorders and personality disorders as risk factors for suicide. Eur Arch Psychiatry Clin Neurosci, 256: 17-27.  Schoenbaum M, Kessler RC, Gilman SE, et al (2014). Predictors of suicide and accident death in the Army Study to Assess Risk and Resilience in Servicemembers (Army STARRS): results from the Army Study to Assess Risk and Resilience in Servicemembers (Army	53215	
Ryding E, Lindstrom M, Traskman-Bendz L (2008). The role of dopamine and serotonin in suicidal behaviour and aggression. Progress in Brain Research, Chapter 15, 172: 307-15.  Sahin HA, Sahin I, Arabaci F (2003). Sociodemographic factors in organophosphate poisonings: a prospective study. Human & Experimental Toxicology, 22: 349-53.  Sakinofsky I (2003). Suicide: the persisting challenge. Can J Psychiatry, 48(5): 289-91.  Sakuta T (1995). A study of murder followed by suicide. Med Law, 14: 141-53.  78683 Sanofi-Aventis Australia (2012). Plaquenil. Product Information.  Santiago JM, Dalen JE (1994). Cholesterol and violent behaviour. Arch Intern Med, 154: 1317-21.  Sareen J, Belik S-L (2009). [Comment] The need for outreach in preventing suicide among young veterans. PLoS Med, 6(3): e1000035. Comment on ID: 53445.  Sareen J, Cox BJ, Afifi TO, et al (2005). Anxiety disorders and risk for suicidal ideation and suicide attempts. Arch Gen Psychiatry, 62: 1249-57.  Sareen J, Cox BJ, Afifi TO, et al (2007). Combat and peacekeeping operations in relation to prevalence of mental disorders and perceived need for mental health care. Arch Gen Psychiatry, 64(7): 843-52.  Schaffer LC, Schaffer CB, Hunter S, et al (2010). Psychiatric reactions to isotretinoin in patients with bipolar disorder. Journal of Affective Disorder, 122(3): 306-8.  Schneider B, Wetterling T, Sargk D, et al (2006). Axis I disorders and personality disorders as risk factors for suicide. Eur Arch Psychiatry Clin Neurosci, 256: 17-27.  Schoenbaum M, Kessler RC, Gilman SE, et al (2014). Predictors of suicide and accident death in the Army Study to Assess Risk and Resilience in Servicemembers (Army Study to Assess Risk and Resilience in Servicemembers (Army	33213	
and serotonin in suicidal behaviour and aggression. Progress in Brain Research, Chapter 15, 172: 307-15.  Sahin HA, Sahin I, Arabaci F (2003). Sociodemographic factors in organophosphate poisonings: a prospective study. Human & Experimental Toxicology, 22: 349-53.  Sakinofsky I (2003). Suicide: the persisting challenge. Can J Psychiatry, 48(5): 289-91.  Sakuta T (1995). A study of murder followed by suicide. Med Law, 14: 141-53.  78683 Sanofi-Aventis Australia (2012). Plaquenil. Product Information.  Santiago JM, Dalen JE (1994). Cholesterol and violent behaviour. Arch Intern Med, 154: 1317-21.  Sareen J, Belik S-L (2009). [Comment] The need for outreach in preventing suicide among young veterans. PLoS Med, 6(3): e1000035. Comment on ID: 53445.  Sareen J, Cox BJ, Affif TO, et al (2005). Anxiety disorders and risk for suicidal ideation and suicide attempts. Arch Gen Psychiatry, 62: 1249-57.  Sareen J, Cox BJ, Affif TO, et al (2007). Combat and peacekeeping operations in relation to prevalence of mental disorders and perceived need for mental health care. Arch Gen Psychiatry, 64(7): 843-52.  Schaffer LC, Schaffer CB, Hunter S, et al (2010). Psychiatric reactions to isotretinoin in patients with bipolar disorder. Journal of Affective Disorder, 122(3): 306-8.  Schneider B, Wetterling T, Sargk D, et al (2006). Axis I disorders and personality disorders as risk factors for suicide. Eur Arch Psychiatry Clin Neurosci, 256: 17-27.  Schoenbaum M, Kessler RC, Gilman SE, et al (2014). Predictors of suicide and accident death in the Army Study to Assess Risk and Resilience in Servicemembers (Army		
Research, Chapter 15, 172: 307-15.  Sahin HA, Sahin I, Arabaci F (2003). Sociodemographic factors in organophosphate poisonings: a prospective study. Human & Experimental Toxicology, 22: 349-53.  Sakinofsky I (2003). Suicide: the persisting challenge. Can J Psychiatry, 48(5): 289-91.  Sakuta T (1995). A study of murder followed by suicide. Med Law, 14: 141-53.  78683 Sanofi-Aventis Australia (2012). Plaquenil. Product Information.  Santiago JM, Dalen JE (1994). Cholesterol and violent behaviour. Arch Intern Med, 154: 1317-21.  Sareen J, Belik S-L (2009). [Comment] The need for outreach in preventing suicide among young veterans. PLoS Med, 6(3): e1000035. Comment on ID: 53445.  Sareen J, Cox BJ, Affif TO, et al (2005). Anxiety disorders and risk for suicidal ideation and suicide attempts. Arch Gen Psychiatry, 62: 1249-57.  Sareen J, Cox BJ, Affif TO, et al (2007). Combat and peacekeeping operations in relation to prevalence of mental disorders and perceived need for mental health care. Arch Gen Psychiatry, 64(7): 843-52.  Schaffer LC, Schaffer CB, Hunter S, et al (2010). Psychiatric reactions to isotretinoin in patients with bipolar disorder. Journal of Affective Disorder, 122(3): 306-8.  Schneider B, Wetterling T, Sargk D, et al (2006). Axis I disorders and personality disorders as risk factors for suicide. Eur Arch Psychiatry Clin Neurosci, 256: 17-27.  Schoenbaum M, Kessler RC, Gilman SE, et al (2014). Predictors of suicide and accident death in the Army Study to Assess Risk and Resilience in Servicemembers (Army STARRS): results from the Army Study to Assess Risk and Resilience in Servicemembers (Army		
Sahin HA, Sahin I, Arabaci F (2003). Sociodemographic factors in organophosphate poisonings: a prospective study. Human & Experimental Toxicology, 22: 349-53.  54029 Sakinofsky I (2003). Suicide: the persisting challenge. Can J Psychiatry, 48(5): 289-91.  52439 Sakuta T (1995). A study of murder followed by suicide. Med Law, 14: 141-53.  78683 Sanofi-Aventis Australia (2012). Plaquenil. Product Information.  Santiago JM, Dalen JE (1994). Cholesterol and violent behaviour. Arch Intern Med, 154: 1317-21.  Sareen J, Belik S-L (2009). [Comment] The need for outreach in preventing suicide among young veterans. PLoS Med, 6(3): e1000035. Comment on ID: 53445.  54734 Sareen J, Cox BJ, Afifi TO, et al (2005). Anxiety disorders and risk for suicidal ideation and suicide attempts. Arch Gen Psychiatry, 62: 1249-57.  Sareen J, Cox BJ, Afifi TO, et al (2007). Combat and peacekeeping operations in relation to prevalence of mental disorders and perceived need for mental health care. Arch Gen Psychiatry, 64(7): 843-52.  Schaffer LC, Schaffer CB, Hunter S, et al (2010). Psychiatric reactions to isotretinoin in patients with bipolar disorder. Journal of Affective Disorder, 122(3): 306-8.  Schneider B, Wetterling T, Sargk D, et al (2006). Axis I disorders and personality disorders as risk factors for suicide. Eur Arch Psychiatry Clin Neurosci, 256: 17-27.  Schoenbaum M, Kessler RC, Gilman SE, et al (2014). Predictors of suicide and accident death in the Army Study to Assess Risk and Resilience in Servicemembers (Army Starks): results from the Army Study to Assess Risk and Resilience in Servicemembers (Army	52965	
organophosphate poisonings: a prospective study. Human & Experimental Toxicology, 22: 349-53.  Sakinofsky I (2003). Suicide: the persisting challenge. Can J Psychiatry, 48(5): 289-91.  Sakuta T (1995). A study of murder followed by suicide. Med Law, 14: 141-53.  78683 Sanofi-Aventis Australia (2012). Plaquenil. Product Information.  Santiago JM, Dalen JE (1994). Cholesterol and violent behaviour. Arch Intern Med, 154: 1317-21.  Sareen J, Belik S-L (2009). [Comment] The need for outreach in preventing suicide among young veterans. PLoS Med, 6(3): e1000035. Comment on ID: 53445.  Sareen J, Cox BJ, Affif TO, et al (2005). Anxiety disorders and risk for suicidal ideation and suicide attempts. Arch Gen Psychiatry, 62: 1249-57.  Sareen J, Cox BJ, Affif TO, et al (2007). Combat and peacekeeping operations in relation to prevalence of mental disorders and perceived need for mental health care. Arch Gen Psychiatry, 64(7): 843-52.  Schaffer LC, Schaffer CB, Hunter S, et al (2010). Psychiatric reactions to isotretinoin in patients with bipolar disorder. Journal of Affective Disorder, 122(3): 306-8.  Schneider B, Wetterling T, Sargk D, et al (2006). Axis I disorders and personality disorders as risk factors for suicide. Eur Arch Psychiatry Clin Neurosci, 256: 17-27.  Schoenbaum M, Kessler RC, Gilman SE, et al (2014). Predictors of suicide and accident death in the Army Study to Assess Risk and Resilience in Servicemembers (Army STARRS): results from the Army Study to Assess Risk and Resilience in Servicemembers (Army		
Experimental Toxicology, 22: 349-53.  Sakinofsky I (2003). Suicide: the persisting challenge. Can J Psychiatry, 48(5): 289-91.  Sakuta T (1995). A study of murder followed by suicide. Med Law, 14: 141-53.  Sanofi-Aventis Australia (2012). Plaquenil. Product Information.  Santiago JM, Dalen JE (1994). Cholesterol and violent behaviour. Arch Intern Med, 154: 1317-21.  Sareen J, Belik S-L (2009). [Comment] The need for outreach in preventing suicide among young veterans. PLoS Med, 6(3): e1000035. Comment on ID: 53445.  Sareen J, Cox BJ, Affif TO, et al (2005). Anxiety disorders and risk for suicidal ideation and suicide attempts. Arch Gen Psychiatry, 62: 1249-57.  Sareen J, Cox BJ, Affif TO, et al (2007). Combat and peacekeeping operations in relation to prevalence of mental disorders and perceived need for mental health care. Arch Gen Psychiatry, 64(7): 843-52.  Schaffer LC, Schaffer CB, Hunter S, et al (2010). Psychiatric reactions to isotretinoin in patients with bipolar disorder. Journal of Affective Disorder, 122(3): 306-8.  Schneider B, Wetterling T, Sargk D, et al (2006). Axis I disorders and personality disorders as risk factors for suicide. Eur Arch Psychiatry Clin Neurosci, 256: 17-27.  Schoenbaum M, Kessler RC, Gilman SE, et al (2014). Predictors of suicide and accident death in the Army Study to Assess Risk and Resilience in Servicemembers (Army STARRS): results from the Army Study to Assess Risk and Resilience in Servicemembers (Army	50000	
Sakinofsky I (2003). Suicide: the persisting challenge. Can J Psychiatry, 48(5): 289-91.  Sakuta T (1995). A study of murder followed by suicide. Med Law, 14: 141-53.  78683 Sanofi-Aventis Australia (2012). Plaquenil. Product Information.  Santiago JM, Dalen JE (1994). Cholesterol and violent behaviour. Arch Intern Med, 154: 1317-21.  Sareen J, Belik S-L (2009). [Comment] The need for outreach in preventing suicide among young veterans. PLoS Med, 6(3): e1000035. Comment on ID: 53445.  Sareen J, Cox BJ, Affif TO, et al (2005). Anxiety disorders and risk for suicidal ideation and suicide attempts. Arch Gen Psychiatry, 62: 1249-57.  Sareen J, Cox BJ, Affif TO, et al (2007). Combat and peacekeeping operations in relation to prevalence of mental disorders and perceived need for mental health care. Arch Gen Psychiatry, 64(7): 843-52.  Schaffer LC, Schaffer CB, Hunter S, et al (2010). Psychiatric reactions to isotretinoin in patients with bipolar disorder. Journal of Affective Disorder, 122(3): 306-8.  Schneider B, Wetterling T, Sargk D, et al (2006). Axis I disorders and personality disorders as risk factors for suicide. Eur Arch Psychiatry Clin Neurosci, 256: 17-27.  Schoenbaum M, Kessler RC, Gilman SE, et al (2014). Predictors of suicide and accident death in the Army Study to Assess Risk and Resilience in Servicemembers (Army STARRS): results from the Army Study to Assess Risk and Resilience in Servicemembers (Army	53996	
48(5): 289-91.  52439  Sakuta T (1995). A study of murder followed by suicide. Med Law, 14: 141-53.  78683  Sanofi-Aventis Australia (2012). Plaquenil. Product Information.  4194  Santiago JM, Dalen JE (1994). Cholesterol and violent behaviour. Arch Intern Med, 154: 1317-21.  Sareen J, Belik S-L (2009). [Comment] The need for outreach in preventing suicide among young veterans. PLoS Med, 6(3): e1000035. Comment on ID: 53445.  Sareen J, Cox BJ, Affif TO, et al (2005). Anxiety disorders and risk for suicidal ideation and suicide attempts. Arch Gen Psychiatry, 62: 1249-57.  Sareen J, Cox BJ, Affif TO, et al (2007). Combat and peacekeeping operations in relation to prevalence of mental disorders and perceived need for mental health care. Arch Gen Psychiatry, 64(7): 843-52.  Schaffer LC, Schaffer CB, Hunter S, et al (2010). Psychiatric reactions to isotretinoin in patients with bipolar disorder. Journal of Affective Disorder, 122(3): 306-8.  Schneider B, Wetterling T, Sargk D, et al (2006). Axis I disorders and personality disorders as risk factors for suicide. Eur Arch Psychiatry Clin Neurosci, 256: 17-27.  Schoenbaum M, Kessler RC, Gilman SE, et al (2014). Predictors of suicide and accident death in the Army Study to Assess Risk and Resilience in Servicemembers (Army STARRS): results from the Army Study to Assess Risk and Resilience in Servicemembers (Army		
Sakuta T (1995). A study of murder followed by suicide. Med Law, 14: 141-53.  78683 Sanofi-Aventis Australia (2012). Plaquenil. Product Information.  4194 Santiago JM, Dalen JE (1994). Cholesterol and violent behaviour. Arch Intern Med, 154: 1317-21.  Sareen J, Belik S-L (2009). [Comment] The need for outreach in preventing suicide among young veterans. PLoS Med, 6(3): e1000035. Comment on ID: 53445.  54734 Sareen J, Cox BJ, Afifi TO, et al (2005). Anxiety disorders and risk for suicidal ideation and suicide attempts. Arch Gen Psychiatry, 62: 1249-57.  Sareen J, Cox BJ, Afifi TO, et al (2007). Combat and peacekeeping operations in relation to prevalence of mental disorders and perceived need for mental health care. Arch Gen Psychiatry, 64(7): 843-52.  Schaffer LC, Schaffer CB, Hunter S, et al (2010). Psychiatric reactions to isotretinoin in patients with bipolar disorder. Journal of Affective Disorder, 122(3): 306-8.  Schneider B, Wetterling T, Sargk D, et al (2006). Axis I disorders and personality disorders as risk factors for suicide. Eur Arch Psychiatry Clin Neurosci, 256: 17-27.  Schoenbaum M, Kessler RC, Gilman SE, et al (2014). Predictors of suicide and accident death in the Army Study to Assess Risk and Resilience in Servicemembers (Army STARRS): results from the Army Study to Assess Risk and Resilience in Servicemembers (Army	54029	
141-53.  78683 Sanofi-Aventis Australia (2012). Plaquenil. Product Information.  4194 Santiago JM, Dalen JE (1994). Cholesterol and violent behaviour. Arch Intern Med, 154: 1317-21.  Sareen J, Belik S-L (2009). [Comment] The need for outreach in preventing suicide among young veterans. PLoS Med, 6(3): e1000035. Comment on ID: 53445.  54734 Sareen J, Cox BJ, Affif TO, et al (2005). Anxiety disorders and risk for suicidal ideation and suicide attempts. Arch Gen Psychiatry, 62: 1249-57.  Sareen J, Cox BJ, Affif TO, et al (2007). Combat and peacekeeping operations in relation to prevalence of mental disorders and perceived need for mental health care. Arch Gen Psychiatry, 64(7): 843-52.  Schaffer LC, Schaffer CB, Hunter S, et al (2010). Psychiatric reactions to isotretinoin in patients with bipolar disorder. Journal of Affective Disorder, 122(3): 306-8.  Schneider B, Wetterling T, Sargk D, et al (2006). Axis I disorders and personality disorders as risk factors for suicide. Eur Arch Psychiatry Clin Neurosci, 256: 17-27.  Schoenbaum M, Kessler RC, Gilman SE, et al (2014). Predictors of suicide and accident death in the Army Study to Assess Risk and Resilience in Servicemembers (Army STARRS): results from the Army Study to Assess Risk and Resilience in Servicemembers (Army		
Santiago JM, Dalen JE (1994). Cholesterol and violent behaviour. Arch Intern Med, 154: 1317-21.  Sareen J, Belik S-L (2009). [Comment] The need for outreach in preventing suicide among young veterans. PLoS Med, 6(3): e1000035. Comment on ID: 53445.  Sareen J, Cox BJ, Afifi TO, et al (2005). Anxiety disorders and risk for suicidal ideation and suicide attempts. Arch Gen Psychiatry, 62: 1249-57.  Sareen J, Cox BJ, Afifi TO, et al (2007). Combat and peacekeeping operations in relation to prevalence of mental disorders and perceived need for mental health care. Arch Gen Psychiatry, 64(7): 843-52.  Schaffer LC, Schaffer CB, Hunter S, et al (2010). Psychiatric reactions to isotretinoin in patients with bipolar disorder. Journal of Affective Disorder, 122(3): 306-8.  Schneider B, Wetterling T, Sargk D, et al (2006). Axis I disorders and personality disorders as risk factors for suicide. Eur Arch Psychiatry Clin Neurosci, 256: 17-27.  Schoenbaum M, Kessler RC, Gilman SE, et al (2014). Predictors of suicide and accident death in the Army Study to Assess Risk and Resilience in Servicemembers (Army STARRS): results from the Army Study to Assess Risk and Resilience in Servicemembers (Army	52439	141-53.
Intern Med, 154: 1317-21.  Sareen J, Belik S-L (2009). [Comment] The need for outreach in preventing suicide among young veterans. PLoS Med, 6(3): e1000035. Comment on ID: 53445.  Sareen J, Cox BJ, Afifi TO, et al (2005). Anxiety disorders and risk for suicidal ideation and suicide attempts. Arch Gen Psychiatry, 62: 1249-57.  Sareen J, Cox BJ, Afifi TO, et al (2007). Combat and peacekeeping operations in relation to prevalence of mental disorders and perceived need for mental health care. Arch Gen Psychiatry, 64(7): 843-52.  Schaffer LC, Schaffer CB, Hunter S, et al (2010). Psychiatric reactions to isotretinoin in patients with bipolar disorder. Journal of Affective Disorder, 122(3): 306-8.  Schneider B, Wetterling T, Sargk D, et al (2006). Axis I disorders and personality disorders as risk factors for suicide. Eur Arch Psychiatry Clin Neurosci, 256: 17-27.  Schoenbaum M, Kessler RC, Gilman SE, et al (2014). Predictors of suicide and accident death in the Army Study to Assess Risk and Resilience in Servicemembers (Army STARRS): results from the Army Study to Assess Risk and Resilience in Servicemembers (Army	78683	
Sareen J, Belik S-L (2009). [Comment] The need for outreach in preventing suicide among young veterans. PLoS Med, 6(3): e1000035. Comment on ID: 53445.  54734  Sareen J, Cox BJ, Afifi TO, et al (2005). Anxiety disorders and risk for suicidal ideation and suicide attempts. Arch Gen Psychiatry, 62: 1249-57.  Sareen J, Cox BJ, Afifi TO, et al (2007). Combat and peacekeeping operations in relation to prevalence of mental disorders and perceived need for mental health care. Arch Gen Psychiatry, 64(7): 843-52.  Schaffer LC, Schaffer CB, Hunter S, et al (2010). Psychiatric reactions to isotretinoin in patients with bipolar disorder. Journal of Affective Disorder, 122(3): 306-8.  Schneider B, Wetterling T, Sargk D, et al (2006). Axis I disorders and personality disorders as risk factors for suicide. Eur Arch Psychiatry Clin Neurosci, 256: 17-27.  Schoenbaum M, Kessler RC, Gilman SE, et al (2014). Predictors of suicide and accident death in the Army Study to Assess Risk and Resilience in Servicemembers (Army STARRS): results from the Army Study to Assess Risk and Resilience in Servicemembers (Army	4194	
53502 preventing suicide among young veterans. PLoS Med, 6(3): e1000035. Comment on ID: 53445.  54734 Sareen J, Cox BJ, Afifi TO, et al (2005). Anxiety disorders and risk for suicidal ideation and suicide attempts. Arch Gen Psychiatry, 62: 1249-57.  Sareen J, Cox BJ, Afifi TO, et al (2007). Combat and peacekeeping operations in relation to prevalence of mental disorders and perceived need for mental health care. Arch Gen Psychiatry, 64(7): 843-52.  Schaffer LC, Schaffer CB, Hunter S, et al (2010). Psychiatric reactions to isotretinoin in patients with bipolar disorder. Journal of Affective Disorder, 122(3): 306-8.  Schneider B, Wetterling T, Sargk D, et al (2006). Axis I disorders and personality disorders as risk factors for suicide. Eur Arch Psychiatry Clin Neurosci, 256: 17-27.  Schoenbaum M, Kessler RC, Gilman SE, et al (2014). Predictors of suicide and accident death in the Army Study to Assess Risk and Resilience in Servicemembers (Army STARRS): results from the Army Study to Assess Risk and Resilience in Servicemembers (Army	4154	
Comment on ID: 53445.  Sareen J, Cox BJ, Afifi TO, et al (2005). Anxiety disorders and risk for suicidal ideation and suicide attempts. Arch Gen Psychiatry, 62: 1249-57.  Sareen J, Cox BJ, Afifi TO, et al (2007). Combat and peacekeeping operations in relation to prevalence of mental disorders and perceived need for mental health care. Arch Gen Psychiatry, 64(7): 843-52.  Schaffer LC, Schaffer CB, Hunter S, et al (2010). Psychiatric reactions to isotretinoin in patients with bipolar disorder. Journal of Affective Disorder, 122(3): 306-8.  Schneider B, Wetterling T, Sargk D, et al (2006). Axis I disorders and personality disorders as risk factors for suicide. Eur Arch Psychiatry Clin Neurosci, 256: 17-27.  Schoenbaum M, Kessler RC, Gilman SE, et al (2014). Predictors of suicide and accident death in the Army Study to Assess Risk and Resilience in Servicemembers (Army Study to Assess Risk and Resilience in Servicemembers (Army		
Sareen J, Cox BJ, Afifi TO, et al (2005). Anxiety disorders and risk for suicidal ideation and suicide attempts. Arch Gen Psychiatry, 62: 1249-57.  Sareen J, Cox BJ, Afifi TO, et al (2007). Combat and peacekeeping operations in relation to prevalence of mental disorders and perceived need for mental health care. Arch Gen Psychiatry, 64(7): 843-52.  Schaffer LC, Schaffer CB, Hunter S, et al (2010). Psychiatric reactions to isotretinoin in patients with bipolar disorder. Journal of Affective Disorder, 122(3): 306-8.  Schneider B, Wetterling T, Sargk D, et al (2006). Axis I disorders and personality disorders as risk factors for suicide. Eur Arch Psychiatry Clin Neurosci, 256: 17-27.  Schoenbaum M, Kessler RC, Gilman SE, et al (2014). Predictors of suicide and accident death in the Army Study to Assess Risk and Resilience in Servicemembers (Army STARRS): results from the Army Study to Assess Risk and Resilience in Servicemembers (Army	53502	
suicidal ideation and suicide attempts. Arch Gen Psychiatry, 62: 1249-57.  Sareen J, Cox BJ, Afifi TO, et al (2007). Combat and peacekeeping operations in relation to prevalence of mental disorders and perceived need for mental health care. Arch Gen Psychiatry, 64(7): 843-52.  Schaffer LC, Schaffer CB, Hunter S, et al (2010). Psychiatric reactions to isotretinoin in patients with bipolar disorder. Journal of Affective Disorder, 122(3): 306-8.  Schneider B, Wetterling T, Sargk D, et al (2006). Axis I disorders and personality disorders as risk factors for suicide. Eur Arch Psychiatry Clin Neurosci, 256: 17-27.  Schoenbaum M, Kessler RC, Gilman SE, et al (2014). Predictors of suicide and accident death in the Army Study to Assess Risk and Resilience in Servicemembers (Army STARRS): results from the Army Study to Assess Risk and Resilience in Servicemembers (Army		
Sareen J, Cox BJ, Afifi TO, et al (2007). Combat and peacekeeping operations in relation to prevalence of mental disorders and perceived need for mental health care. Arch Gen Psychiatry, 64(7): 843-52.  Schaffer LC, Schaffer CB, Hunter S, et al (2010). Psychiatric reactions to isotretinoin in patients with bipolar disorder. Journal of Affective Disorder, 122(3): 306-8.  Schneider B, Wetterling T, Sargk D, et al (2006). Axis I disorders and personality disorders as risk factors for suicide. Eur Arch Psychiatry Clin Neurosci, 256: 17-27.  Schoenbaum M, Kessler RC, Gilman SE, et al (2014). Predictors of suicide and accident death in the Army Study to Assess Risk and Resilience in Servicemembers (Army STARRS): results from the Army Study to Assess Risk and Resilience in Servicemembers (Army	54734	
operations in relation to prevalence of mental disorders and perceived need for mental health care. Arch Gen Psychiatry, 64(7): 843-52.  Schaffer LC, Schaffer CB, Hunter S, et al (2010). Psychiatric reactions to isotretinoin in patients with bipolar disorder. Journal of Affective Disorder, 122(3): 306-8.  Schneider B, Wetterling T, Sargk D, et al (2006). Axis I disorders and personality disorders as risk factors for suicide. Eur Arch Psychiatry Clin Neurosci, 256: 17-27.  Schoenbaum M, Kessler RC, Gilman SE, et al (2014). Predictors of suicide and accident death in the Army Study to Assess Risk and Resilience in Servicemembers (Army STARRS): results from the Army Study to Assess Risk and Resilience in Servicemembers (Army		
need for mental health care. Arch Gen Psychiatry, 64(7): 843-52.  Schaffer LC, Schaffer CB, Hunter S, et al (2010). Psychiatric reactions to isotretinoin in patients with bipolar disorder. Journal of Affective Disorder, 122(3): 306-8.  Schneider B, Wetterling T, Sargk D, et al (2006). Axis I disorders and personality disorders as risk factors for suicide. Eur Arch Psychiatry Clin Neurosci, 256: 17-27.  Schoenbaum M, Kessler RC, Gilman SE, et al (2014). Predictors of suicide and accident death in the Army Study to Assess Risk and Resilience in Servicemembers (Army STARRS): results from the Army Study to Assess Risk and Resilience in Servicemembers (Army	52004	
Schaffer LC, Schaffer CB, Hunter S, et al (2010). Psychiatric reactions to isotretinoin in patients with bipolar disorder. Journal of Affective Disorder, 122(3): 306-8.  Schneider B, Wetterling T, Sargk D, et al (2006). Axis I disorders and personality disorders as risk factors for suicide. Eur Arch Psychiatry Clin Neurosci, 256: 17-27.  Schoenbaum M, Kessler RC, Gilman SE, et al (2014). Predictors of suicide and accident death in the Army Study to Assess Risk and Resilience in Servicemembers (Army STARRS): results from the Army Study to Assess Risk and Resilience in Servicemembers (Army	52094	
isotretinoin in patients with bipolar disorder. Journal of Affective Disorder, 122(3): 306-8.  Schneider B, Wetterling T, Sargk D, et al (2006). Axis I disorders and personality disorders as risk factors for suicide. Eur Arch Psychiatry Clin Neurosci, 256: 17-27.  Schoenbaum M, Kessler RC, Gilman SE, et al (2014). Predictors of suicide and accident death in the Army Study to Assess Risk and Resilience in Servicemembers (Army STARRS): results from the Army Study to Assess Risk and Resilience in Servicemembers (Army		
122(3): 306-8.  Schneider B, Wetterling T, Sargk D, et al (2006). Axis I disorders and personality disorders as risk factors for suicide. Eur Arch Psychiatry Clin Neurosci, 256: 17-27.  Schoenbaum M, Kessler RC, Gilman SE, et al (2014). Predictors of suicide and accident death in the Army Study to Assess Risk and Resilience in Servicemembers (Army STARRS): results from the Army Study to Assess Risk and Resilience in Servicemembers (Army	77961	
Schneider B, Wetterling T, Sargk D, et al (2006). Axis I disorders and personality disorders as risk factors for suicide. Eur Arch Psychiatry Clin Neurosci, 256: 17-27.  Schoenbaum M, Kessler RC, Gilman SE, et al (2014). Predictors of suicide and accident death in the Army Study to Assess Risk and Resilience in Servicemembers (Army STARRS): results from the Army Study to Assess Risk and Resilience in Servicemembers (Army		· ·
49905 personality disorders as risk factors for suicide. Eur Arch Psychiatry Clin Neurosci, 256: 17-27.  Schoenbaum M, Kessler RC, Gilman SE, et al (2014). Predictors of suicide and accident death in the Army Study to Assess Risk and Resilience in Servicemembers (Army STARRS): results from the Army Study to Assess Risk and Resilience in Servicemembers (Army	49905	
Neurosci, 256: 17-27.  Schoenbaum M, Kessler RC, Gilman SE, et al (2014). Predictors of suicide and accident death in the Army Study to Assess Risk and Resilience in Servicemembers (Army STARRS): results from the Army Study to Assess Risk and Resilience in Servicemembers (Army		
Schoenbaum M, Kessler RC, Gilman SE, et al (2014). Predictors of suicide and accident death in the Army Study to Assess Risk and Resilience in Servicemembers (Army STARRS): results from the Army Study to Assess Risk and Resilience in Servicemembers (Army		
suicide and accident death in the Army Study to Assess Risk and Resilience in Servicemembers (Army STARRS): results from the Army Study to Assess Risk and Resilience in Servicemembers (Army		
Resilience in Servicemembers (Army STARRS): results from the Army Study to Assess Risk and Resilience in Servicemembers (Army		
Study to Assess Risk and Resilience in Servicemembers (Army	83208	
STARRS). JAMA Psychiatry, 71(5): 493-503.		Study to Assess Risk and Resilience in Servicemembers (Army
		STARRS). JAMA Psychiatry, 71(5): 493-503.

	Schreiber J, Culpepper L (2016). Suicidal ideation and behavior in adults.  Retrieved 14 March 2016, from
77719	http://www.uptodate.com/contents/suicidal-ideation-and-behavior-in-adults
54037	Scoville SL, Gubata ME, Potter RN, et al (2007). Deaths attributed to suicide among enlisted U.S. armed forces recruits, 1980-2004. Military Medicine, 172(10): 1024-31.
77840	Scudder L, Seymour S, Torjusen E, et al (2015). Montelukast's underrecognized adverse drug events. Retrieved 24 March 2016, from http://www.medscape.com/viewarticle/840302#vp_1
11040	Senate Standing Committee on Foreign Affairs, Defence and Trade (1994). Facing the Future Together. Sexual Harassment in the Australian Defence Force, Chapter 1: 1-2, 32. Commonwealth of Australia, Parliament House, Canberra.
43604	Shalev AY, Tuval-Mashiach R, Hadar H (2004). Posttraumatic Stress Disorder as a result of mass trauma. Journal of Clinical Psychiatry, 65(S1): 4-10.
77798	Sharma T, Guski LS, Freund N, et al (2016). Suicidality and aggression during antidepressant treatment: systematic review and meta-analyses based on clinical study reports. BMJ, 352: i65.
52661	Shaw KR, Sheehan KH, Fernandez RC (1987). Suicide in children and adolescents. Adv Pediatr, 34: 313-34.
52457	Shelton AJ, Atkinson J, Risser JM, et al (2006). The prevalence of suicidal behaviours in a group of HIV-positive men. AIDS Care, 18(6): 574-6.
79959	Shen YC, Cunha JM, Williams TV (2016). Time-varying associations of suicide with deployments, mental health conditions, and stressful life events among current and former US military personnel: a retrospective multivariate analysis. [Epub ahead of print]. Lancet Psychiatry.
52678	Sherer DM, Schenker JG (1989). Accidental injury during pregnancy. Obstet Gynecol Surv, 44(5): 330-8.
52444	Sherr L, Lampe F, Fisher M, et al (2008). Suicidal ideation in UK HIV clinic attenders. AIDS, 22: 1651-8.
54680	Shneidman E (2005). Prediction of suicide revisited: a brief methodological note. Suicide & Life-Threatening Behaviour, 35(1): 1-2.
76850	Siamouli M, Samara M, Fountoulakis KN (2014). Is antiepileptic-induced suicidality a data-based class effect or an exaggeration? A comment on the literature. Harv Rev Psychiatry, 22(6): 379-81.
54050	Siegert RJ, Abernethy DA (2005). Depression in multiple sclerosis: a review. J Neurol Neurosurg Psychiatry, 76: 469-75.
52453	Silenzio VM, Pena JB, Duberstein PR, et al (2007). Sexual orientation and risk factors for suicidal ideation and suicide attempts among adolescents and young adults. Am J Public Health, 97(11): 2017-9.
54901	Silverman MM, Berman AL, Sanddal ND, et al (2007). Rebuilding the tower of Babel: a revised nomenclature for the study of suicide and suicidal behaviors. Part 1: background, rationale, and methodology. What, 37(3): 248-63.
31718	Sim M, Abramson M, Forbes A, et al (2003). Psychological Health. Australian Gulf War Veterans' Health Study, Vol 2 Chapter 11: 268-71. Commonwealth of Australia.
77791	Simon G (2015). Effect of antidepressants on suicide risk in adults Retrieved 17 March 2016, from www.uptodate.com/contents/effect-of-antidepressants-on-suicide-risk-in-adults
48439	Simon RI (2008). Suicide. Textbook of Psychiatry, 5th Edition, Part VI, Chapter 43. American Psychiatric Publishing, Inc.

-	<del>-</del>
77768	Singhal A, Ross J, Seminog O, et al (2014). Risk of self-harm and suicide in people with specific psychiatric and physical disorders: comparisons between disorders using English national record linkage. J Roy Soc Med, 107(5): 194-204.
77617	Sinyor M, Cheung AH (2015). Antidepressants and risk of suicide. BMJ, 350: h783.
84408	Skogstad L, Fjetland AM, Ekeberg O (2015). Exposure and posttraumatic stress symptoms among first responders working in proximity to the terror sites in Norway on July 22, 2011 - a cross-sectional study. Scand J Trauma Resusc Emerg Med, 23(23): 1-9.
83742	Skogstad L, Heir T, Hauff E, et al (2016). Post-traumatic stress among rescue workers after terror attacks in Norway. Occup Med, 66(7): 528-35.
79640	Skopp NA, Zhang Y, Smolenski DJ, et al (2016). Risk factors for self-directed violence in US soldiers: a case-control study. Psychiatry Res, 245: 194-9.
54030	Smith JP, Riley TR, Bingaman S, et al (2004). Amantadine therapy for chronic hepatitis C: a dose escalation study. Am J Gastroenterol, 99: 1099-104.
54002	Smith MT, Edwards RR, Robinson RC, et al (2004). Suicidal ideation, plans, and attempts in chronic pain patients: factors associated with increased risk. Pain, 111: 201-8.
77483	Song HR, Woo YS, Wang H-R, et al (2014). How does antiepileptic drug induce suicidality? A case associated with levitracetam use. Gen Hosp Psychiatry, 36(3): 360e1-360e2.
77720	Soreff S (2016). Suicide. Retrieved 14 March 2016, from http://emedicine.medscape.com/article/2013085-overview#a3
54295	Sorensen HT, Mellemkjaer L, Olsen JH (2001). Risk of suicide in users of b-adrenoceptor blockers, calcium channel blockers and angiotensin converting enzyme inhibitors. Br J Clin Pharmacol, 52: 313-8.
54007	Spiegel B, Schoenfelds P, Naliboff B (2007). Systematic review: the prevalence of suicidal behaviour in patients with chronic abdominal pain and irritable bowel syndrome. Aliment Pharmacol Ther, 26: 183-93.
28553	Spooner MH (2002). [Comment] Suicide claiming more British Falkland veterans than fighting did. CMAJ, 166(11): 1453.
77799	Sriretnakumar V, Huang E, Muller D (2015). Pharmacogenetics of clozapine treatment response and side-effects in schizophrenia: an update. Expert Opin Drug Metab Toxicol, 11(11): 1709-31.
83209	Stack S, Scourfield J (2015). Recency of divorce, depression, and suicide risk. Journal of Family Issues, 36(6): 695-715.
54202	Stallones L (2006). Suicide and potential occupational exposure to pesticides, Colorado 1990-1999. J Agromed, 11(3/4): 107-12.
54735	Stanley B, Gameroff MJ, Michalsen V, et al (2001). Are suicide attempters who self-mutilate a unique population? Am J Psychiatry, 158: 427-32.
53467	Starcevic V, Bogojevic G, Marinkovic J, et al (1999). Axis I and axis II comorbidity in panic / agoraphobic patients with and without suicidal ideation. Psychiatr Res, 88: 153-61.
53990	Stark C, Belbin A, Gibbs D, et al (2006). Male suicide and occupation in Scotland. Health Statistics Quarterly, 29: 26-9.
53992	Stark C, Gibbs D, Hopkins P, et al (2006). Suicide in farmers in Scotland. Rural and Remote Health, 6: 509.
79784	Stein DJ, Chiu WT, Hwang I, et al (2010). Cross-national analysis of the association between traumatic events and suicidal behavior: findings from the WHO world mental health surveys. PLoS One, 5(5): e10574.

77844	Stein MB, Kessler RC, Heeringa SG, et al (2015). Prospective longitudinal evaluation of the effect of deployment-acquired traumatic brain injury on posttraumatic stress and related disorders: results from the army study to assess risk and resilience in servicemembers (army STARRS). Am J Psychiatry, 172(11): 1101-11.
1771	Stenager EN, Stenager E, Jensen K (1994). Attempted suicide, depression and physical diseases: a 1-year follow-up study. Psychother Psychosom, 61: 65-73.
53676	Sterud T, Hem E, Lau B, et al (2008). Suicidal ideation and suicide attempts in a nationwide sample of operational Norwegian ambulance personnel. J Occup Health, 50: 406-14.
85930	Stetz MC, Wiederhold BK, Wildzunas RM, et al (2006). The usefulness of virtual reality stress inoculation training for military medical females: A pilot study. Annual Review of CyberTherapy and Telemedicine, 4: 51-8.
86471	Stewart SH, Mitchell TL, Wright KD, et al (2004). The relations of PTSD symptoms to alcohol use and coping drinking in volunteers who responded to the Swissair Flight 111 airline disaster. J Anxiety Dis, 18(1): 51-68.
54025	Stone M, Laughren T, Jones ML, et al (2009). Risk of suicidality in clinical trials of antidepressants in adults: analysis of proprietary data submitted to US Food & Drug Administration. BMJ, 339: b2280.
53453	Strauss JL, Calhoun PS, Marx CE, et al (2006). Comorbid posttraumatic stress disorder is associated with suicidality in male veterans with schizophrenia or schizoaffective disorder. Schizophrenia Research, 84: 165-9.
79678	Substance Abuse and Mental Health Services Administration (US) (2014). Trauma-informed care in behavioral health services. Treatment improvement protocol (TIP) Series, No. 57. HHS Publication No. (SMA) 13-4801, Chapter 3 - Understanding the impact of trauma. US Department of Health and Human Services. Center for Substance Abuse Treatment (US) Rockville (MD).
77808	Sun Y, Lin C-C, Lu C-J, et al (2016). Association between zolpidem and suicide: a nationwide population-based case-control study. Mayo Clin Proc, 91(3): 308-15.
77959	Sundstrom A, Alfredsson L, Siolin-Forsberg G, et al (2010). Association of suicide attempts with acne and treatment with isotretinoin: retrospective Swedish cohort study. BMJ, 341: c5812.
54045	Sunnqvist C, Westrin A, Traskman-Bendz L (2008). Suicide attempters: biological stressmarkers and adverse life events. Eur Arch Psychiatry Clin Neurosci, 258: 456-62.
1772	Suokas J, Jönnqvist J (1995). Suicide attempts in which alcohol is involved: a special group in general hospital emergency rooms. Acta Psychiatr Scand, 91: 36-40.
83741	Surgenor LJ, Snell DL, Dorahy MJ (2015). Posttraumatic stress symptoms in police staff 12-18 months after the Canterbury earthquakes. J Trauma Stress, 28(2): 162-6.
78114	Suzuki J, Dekker MA, Valenti ES, et al (2015). Toxicities associated with NBOMe ingestion, a novel class of potent hallucinogens: a review of the literature. Psychosomatics, 56(2): 129-39.
78113	Suzuki J, Poklis JL, Poklis A (2014). "My friend said it was good LSD". A suicide attempt following analytically confirmed 251-NBOMe ingestion. J Psychoactive Drugs, 46(5): 379-82.
52447	Swahn MH, Simon TR, Hertz MF, et al (2008). Linking dating violence, peer violence, and suicidal behaviors among high-risk youth. Am J Prev Med, 34(1): 30-8.
77695	Swanson SA, Colman I (2013). Association between exposure to suicide and suicidality outcomes in youth. CMAJ, 185(10): 870-7.

1773	Takei N (1994). [Comment] Low serum cholesterol and suicide attempts. Br J Psychiatry, 164(5): 702-3.
53890	Tandt H, Audenaert K, van Heeringen C (2009). SSRIs (selective serotonin reuptake inhibitors) and suicidality in adults, adolescents and children. Tijdschr Psychiatr, 51(6): 387-93. [Abstract]
54004	Tang NK, Crane C (2006). Suicidality in chronic pain: a review of the prevalence, risk factors and psychological links. Psychol Med, 36: 575-86.
54269	Taylor DJ, Lichstein KL, Durrence HH (2003). Insomnia as a health risk factor. Behav Sleep Med, 1(4): 227-47.
1774	Taylor SJ (1994). Debt and deliberate self-poisoning. Br J Psychiatry, 164: 111-4.
54456	Terao T, Soya A, Brunner J, et al (2003). [Comments] Cholesterol, essential fatty acids, and suicide. Pharmacopsychiatry, 36: 83-5. Comments on ID: 54455.
77623	Termorshuizen F, Palmen SJM, Heerdink ER (2015). Suicide behavior before and after the start with antidepressants: a high persistent risk in the first month of treatment among the young. Int J Neuropsychopharmacol, 1-10.
54736	Thacore VR, Varma SL (2000). A study of suicides in Ballarat, Victoria, Australia. Crisis, 21(1): 26-30.
53989	Thomas HV, Lewis G, Thomas D R, et al (2003). Mental health of British farmers. Occup Environ Med, 60: 181-6.
77563	Thomas KH, Martin RM, Davies NM, et al (2013). Smoking cessation treatment and risk of depression, suicide, and self harm in the clinical practice research datalink: prospective cohort study. BMJ, 347: I5704.
77565	Thomas KH, Martin RM, Knipe DW, et al (2015). Risk of neuropsychiatric adverse events associated with varenicline: systematic review and meta-analysis. BMJ, 350: h1109.
77564	Thomas KH, Martin RM, Potokar J, et al (2014). Reporting of drug induced depression and fatal and non-fatal suicidal behaviour in the UK from 1998-2011. BMC Pharmacology and Toxicology, 15: 54.
53459	Thompson R, Kane V, Cook JM, et al (2006). Suicidal ideation in veterans receiving treatment for opiate dependence. Journal of Psychoactive Drugs, 38(2): 149-56.
53462	Thompson R, Katz IR, Kane VR, et al (2002). Cause of death in veterans receiving general medical and mental health care. J Nerv Ment Dis, 190(11): 789-92.
53457	Thoresen S, Mehlum L (2008). Traumatic stress and suicidal ideation in Norwegian male peacekeepers. J Nerv Ment Dis, 196(11): 814-21.
31740	Thoresen S, Mehlum L, Moller B (2003). Suicide in peacekeepers: a cohort study of mortality from suicide in 22,275 Norwegian veterans form international peacekeeping operations. Soc Psychiatry Psychiatr Epidemiol, 38: 605-10.
53686	Thoresen S, Mehlum L, Roysamb E, et al (2006). Risk factors for completed suicide in veterans of peacekeeping: repatriation, negative life events, and marital status. Arch Suicide Res, 10: 353-63.
86469	Thormar SB, Gersons BPR, Juen B, et al (2013). Organizational factors and mental health in community volunteers. The role of exposure, preparation, training, tasks assigned, and support. Anxiety, Stress, & Coping, 26(6): 624-42.
85883	Tierens M, Bal S, Crombez G, et al (2012). Differences in posttraumatic stress reactions between witnesses and direct victims of motor vehicle accidents. Journal of Traumatic Stress, 25(3): 280-7.
53452	Tiet QQ, Finney JW, Moos RH (2006). Recent sexual abuse, physical abuse, and suicide attempts among male veterans seeking psychiatric treatment. Psychiatric Services, 57(1): 107-13.

77935	Tindle HA, Omalu B, Courcoulas A, et al (2010). Risk of suicide after long term follow-up from bariatric surgery. Am J Med, 123(11): 1036-42.
77793	Toffol E, Hatonen T, Tanskanen A, et al (2015). Lithium is associated with decrease in all-cause and suicide mortality in high-risk bipolar patients: A nationwide registry-based prospective cohort study. J Affect
	Disord, 183: 159-65.  Torelli P, Manzoni GC (2003). Pain and behaviour in cluster headache. A
54829	prospective study and review of the literature. Functional Neurology, 18(4): 205-10.
79755	Tran LD, Grant D, Aydin M, et al (2016). California veterans receive inadequate treatment to address their mental health needs. Am J Med Res, 3(2): 126-40.
52470	Tran Thi Thanh H, Tran NT, Jiang G, et al (2006). Life time suicidal thoughts in an urban community in Hanoi, Vietnam. BMC Public Health, 6: 76.
77845	Truven Health Analytics Inc (2016). Levofloxacin: adverse effects Retrieved 24 March 2016, from http://www.micromedexsolutions.com
54768	Tsoh J, Chiu HFK, Duberstein PR, et al (2005). Attempted suicide in elderly Chinese persons. Am J Geriatr Psychiatry, 13(7): 562-71.
53454	Turner AP, Williams RM, Bowen JD, et al (2006). Suicidal ideation in multiple sclerosis. Arch Phys Med Rehabil, 87: 1073-8.
77369	U.S. Department of Health and Human Services (2012). National strategy for suicide prevention: goals and objectives for action. Appendix D: Groups with increased suicide risk. HHS, 101-30. Washington DC US Government Printing Office.
86460	Ursano RJ (1990). The nature of a traumatic stressor: Handling dead bodies. The Journal of Nervous and Mental Disease, 178(6): 396-8.
79642	Ursano RJ, Kessler RC, Stein MB, et al (2016). Risk factors, methods, and timing of suicide attempts among US army soldiers. JAMA Psychiatry, 73(7): 741-9.
52442	van Bergen DD, Smit JH, van Balkom AJ, et al (2008). Suicidal ideation in ethnic minority and majority adolescents in Utrecht, The Netherlands. Crisis, 29(2): 202-8.
49904	van der Wal MF, de Wit CAM, Hirasing RA (2003). Psychosocial health among young victims and offenders of direct and indirect bullying. Pediatrics, 111(6): 1312-7.
77570	van Geel M, Vedder P, Tanilon J (2014). Relationship between peer victimization, cyberbullying, and suicide in children and adolescents: a meta-analysis. JAMA Pediatr, 168(5): 435-42.
54028	van Heeringen K (2003). The neurobiology of suicide and suicidality. Can J Psychiatry, 48(5): 292-300.
54571	Vander Stoep A, Weiss NS, McKnight B, et al (2002). Which measure of adolescent psychiatric disorder - diagnosis, number of symptoms, or adaptive functioning - best predicts adverse young adult outcomes? J Epidemiol Community Health, 56: 56-95.
52455	Vanderwerker LC, Chen JH, Charpentier P, et al (2007). Differences in risk factors for suicidality between African American and white patients vulnerable to suicide. Suicide & Life-Threatening Behaviour, 37(1): 1-9.
4196	Vasan RS (1992). Lowering cholesterol and death due to accidents, suicides: Unresolved issues. Arch Intern Med, 152: 414.
54000	Vickers A, Wright M (2002). [Comments] Widespread body pain and mortality. BMJ, 324: 300.
54048	Voaklander DC, Rowe BH, Dryden DM, et al (2008). Medical illness, medication use and suicide in seniors: a population-based case-control study. J Epidemiol Community Health, 62: 138-46.

70941	Vogt D, Vaughn R, Glickman ME, et al (2011). Gender differences in combat-related stressors and their association with postdeployment mental health in a nationally representative sample of U.S. OEF/OIF veterans. J Abnorm Psychol, 120(4): 797-806.
54284	Voracek M, Loibl LM (2007). Genetics of suicide: a systematic review of twin studies. Wien Klin Wochenschr, 119(15-16): 463-75.
54723	Waern M, Rubenowitz E, Runeson B, et al (2002). Burden of illness and suicide in elderly people: case-control study. BMJ, 324: 1355.
54722	Waern M, Rubenowitz E, Wilhelmson K (2003). Predictors of suicide in the old elderly. Gerontology, 49: 328-34.
77371	Wagman Borowsky I, Taliaferro LA, McMorris BJ (2013). Suicidal thinking and behavior among youth involved in verbal and social bullying: risk and protective factors. J Adolesc Health, 53: S4-S12.
52682	Walker RL, Obasi EM, Wingate LR, et al (2008). An empirical investigation of acculturative stress and ethnic identity as moderators for depression and suicidal ideation in college students. Cultur Divers Ethnic Minor Psychol, 14(1): 75-82.
78020	Walsh DS, Eamsila C, Sasiprapha T, et al (2004). Efficacy of monthly tafenoquine for prophylaxis of plasmodium vivax and multidrug-resistant p. falciparum malaria. J Infect Dis, 190(8): 1456-63.
78023	Walsh DS, Looareesuwan S, Wilairantana P, et al (1999). Randomized dose-ranging study of the safety and efficacy or WR238605 (tafenoquine) in the prevention of relapse of plasmodium vivax malaria in Thailand. J Infect Dis, 180(4): 1282-7.
54036	Walsh E, Eggert LL (2007). Suicide risk and protective factors among youth experiencing school difficulties. Int J Ment Health Nurs, 16: 349-59.
54204	Walsh Sm, Sage RA (2002). Depression and chronic diabetic foot disability. A case report of suicide. Clin Podiatr Med Surg, 19: 493-508.
83740	Wang H, Jin H, Nunnink SE, et al (2011). Identification of post traumatic stress disorder and risk factors in military first responders 6 months after Wen Chuan earthquake in China. J Affect Disord, 130(1-2): 213-9.
54769	Wang S-J, Juang K-D, Fuh J-L, Lu S-R (2007). Psychiatric comorbidity and suicide risk in adolescents with chronic daily headache. Neurology, 68: 1468-73.
53446	Warner J, McKeown E, Griffin M, et al (2004). Rates and predictors of mental illness in gay men, lesbians and bisexual men and women. Br J Psychiatry, 185: 479-85.
54215	Warrington TP, Bostwick JM (2006). Psychiatric adverse effects of corticosteroids. Mayo Clin Proc, 81(10): 1361-7.
4176	Warshaw MG, Massion AO, Peterson LG, et al (1995). Suicidal behaviour in patients with panic disorder: retrospective and prospective data. Journal of Affective Disorders, 34: 235-47.
1775	Wasserman D, Spellerberg S (1989). Attempted-suicide trends in Stockholm County, Sweden, 1975-1985. Acta Psychiatr Scand, 81: 190-6.
77567	Weiner J, Richmond TS, Conigllaro J, et al (2011). Military veteran mortality following a survived suicide attempt. BMC Public Health, 11: 374.
75208	Weinke T, Trautmann M, Held T, et al (1991). Neuropsychiatric side effects after the use of mefloquine. Am J Trop Med Hyg, 45(1): 86-91.
52667	Weissman MM, Klerman GL (1977). Sex differences and the epidemiology of depression. Arch Gen Psychiatry, 34: 98-111.
77259	Wells TS, Smith TC, Smith B (2006). Mefloquine use and hospitalizations among US service members, 2002-2004. Am J Trop Med Hyg, 74(5): 744-749.
85890	Werner EE (2012). Children and war: Risk, resilience, and recovery. Development and Psychopathology, 24(2): 553-8.

	THE CO. D. LEWIS CO. L. (2000) 14 A 14 A 14
00470	West C, Bernard B, Mueller C, et al (2008). Mental health outcomes in
86472	police personnel after Hurricane Katrina. J Occup Environ Med, 50(6): 689-95.
	WHO (2014). Preventing suicide: a global imperative. World Health
77566	Organization.
77626	Wijlaars LPMM, Nazareth I, Whitaker HJ, et al (2013). Suicide-related
	events in young people following prescription of SSRIs and other
	antidepressants: a self-controlled case series analysis. BMJ Open, 3:
	e003247.
	Wilcox HC, Conner KR, Caine ED (2004). Association of alcohol and
54021	drug use disorders and completed suicide: an empirical review of cohort
	studies. Drug and Alcohol Dependence, 76S: S11-9.
78396	Wilcox HC, Kuramoto SJ, Linchtenstein P, et al (2010). Psychiatric
	morbidity, violent crime, and suicide among children and adolescents
	exposed to parental death. J Am Acad Child Adolesc Psychiatry, 49(5): 514-23.
	Williams JM, Pollock LR (2000). The psychology of suicidal behaviour.
53216	The International Handbook of Suicide and Attempted Suicide, Chapter
00210	5: 79-93. John Wiley & Sons, New York, US.
	Wilson LC (2015). A systematic review of probable posttraumatic stress
84405	disorder in first responders following man-made mass violence.
	Psychiatry Res, 229: 21-6.
	Windle M (1993). Characteristics of alcoholics who attempted suicide: co-
1776	occurring disorders and personality differences with a sample of male
	Vietnam era veterans. J Stud Alcohol, 55(55): 571-7.
77611	Wise J (2016). Antidepressants may double risk of suicide and
	aggression in children, study finds. BMJ, 352: i545. Witteveen AB, Bramsen I, Twisk WR, et al (2007). Psychological distress
69868	of rescue workers eight and one-half years after professional involvement
03000	in the Amsterdam air disaster. J Nerv Ment Dis, 195(1): 31-40.
	Wojnar M, Ilgen MA, Wojnar J, et al (2009). Sleep problems and
52684	suicidality in the National Comorbidity Survey Replication. J Psychiat
	Res, 43: 526-31.
31739	Wong A, Escobar M, Lesate A, et al (2001). Are UN Peacekeepers at risk
01700	of suicide? Suicide and Life-Threatening Behavior, 31: 103-12.
52452	Wong JP, Stewart SM, Claassen C, et al (2008). Repeat suicide attempts
	in Hong Kong community adolescents. Soc Sci Med, 66: 232-41.
54770	Woolley SB, Fredman L, Goethe JW, et al (2008). Headache complaints and the risk of suicidal thoughts or behaviors. J Nerv Ment Dis, 196(11):
34770	822-8.
00500	Wooltorton E (2003). Accutane (isotretinoin) and psychiatric adverse
38500	effects. JAMC, 168: 66.
77368	World Health Organization (2014). Preventing suicide: a global
11300	imperative. WHO. Geneva, Switzerland.
53500	Wortzel HS, Binswanger IA, Anderson A, et al (2009). Suicide among
30000	incarcerated veterans. J Am Acad Psychiatry Law, 37: 82-91.
54691	Wyder M, De Leo D (2007). Behind impulsive suicide attempts:
	indications from a community study. J Affect Disord, 104: 167-73.  Wysowski DK, Gross TP (1990). Deaths due to accidents and violence in
4197	two recent trials of cholesterol-lowering drugs. Arch Intern Med, 150:
4131	2169-72.
83211	Xiao J, Guan S, Ge H, et al (2017). The impact of changes in work
	stressors and coping resources on the risk of new-onset suicide ideation
	among Chinese petroleum industry workers. J Psychiatry Res, 88: 1-8.
	Yamasaki A, Chinami M, Morgenthaler S, et al (2004). Enterprise failures
54729	correlate positively with suicide rate for both sexes in Japan.
	Psychological Reports, 95: 914-20.

54293	Yen S, Pagano ME, Shea MT, et al (2005). Recent life events preceding suicide attempts in a personality disorder sample: findings from the Collaborative Longitudinal Personality Disorders Study. J Consult Clin Psychol, 73(1): 99-105.
49800	Yen S, Shea MT, Pagano M, et al (2003). Axis I and axis II disorders as predictors of prospective suicide attempts: findings from the collaborative longitudinal personality disorders study. J Abnorm Psychol, 112(3): 375-81. [Erratum attached].
77568	Yi S-W, Hong J-S (2015). Depressive symptoms and other risk factors predicting suicide in middle-aged men: a prospective cohort study among Korean Vietnam war veterans. PeerJ, 3: e1071.
52968	Young AH (2009). Invited commentary on Lithium levels in drinking water and risk of suicide. The British Journal of Psychiatry, 194(5): 466. Retrieved 29 May 2009, from http://bjp.rcpsych.org/cgi/content/full/194/5/466
78115	Zarrabi H, Khalkhali M, Hamidi A, et al (2016). Clinical features, course and treatment of methamphetamine-induced psychosis in psychiatric inpatients. BMC Psychiatry, 16: 44.
77473	Zerach G, Levi-Belz Y, Solomon Z (2014). Trajectories of suicidal ideation and posttraumatic stress symptoms among former prisoners of war: a 17-year longitudinal study. J Psychiatr Res, 49: 83-9.
77599	Zerwas S, Tidselbak Larsen J, Petersen L, et al (2015). The incidence of eating disorders in a Danish register study: associations with suicide risk and mortality. J Psychiatry Res, 65: 16-22.
77759	Zubrick SR, Hafekost J, Johnson SE, et al (2016). Suicidal behaviours: prevalence estimates from the second Australian child and adolescent survey of mental health and wellbeing. Aust N Z J Psychiatry, [Epub ahead of print].